arXiv:1207.6296v2 [math.CO] 27 Jul 2012

The diameters of associahedra

Lionel Pournin

July 27, 2012

Abstract It is proven here, using combinatorial arguments, that the diameter of the d-dimensional associahedron is $2d - 4$ when d is greater than 9.

Keywords Associahedra · flip-graphs · triangulations · polygons · secondary polytopes · rotation distance · binary trees · associativity

Mathematics Subject Classification (2000) 51M20 · 05E45

1 Introduction

The history of associahedra can be traced back to when James Stasheff introduced their face complexes [18], and used them to study homotopy associative H-spaces. They were later constructed as convex polytopes by Mark Haiman in an unpublished manuscript [11], and then by Carl Lee [13]. In the last fifty years, associahedra have attracted much attention, not only due to their importance in algebraic topology, but also because of their connection with a number of combinatorial objects such as triangulations of convex polygons [17,13], binary trees [17,4], and bracketed expressions [12]. Their diameters are especially relevant to operations that can be carried out to transform any of these objects into other objects of the same type: flips in triangulations, rotations in binary trees, and associativity in bracketed expressions.

About twenty-five years ago, Daniel Sleator, Robert Tarjan, and William Thurston proved the following result [17]:

Theorem 1 *When d is large enough, the d-dimensional associahedron has diameter $2d - 4$.*

L. Pournin
EFREI, 30-32 avenue de la République, 94800 Villejuif, France
E-mail: lionel.pournin@efrei.fr
EPFL - École Polytechnique Fédérale de Lausanne, 1015 Lausanne, Switzerland
E-mail: lionel.pournin@epfl.ch

Note that this theorem is originally stated in [17] using the formalism of triangulations and flips, and does not explicitly mentions associahedra. In fact, the connection between associahedra and triangulations of convex polygons was discovered independently at about the same time by Carl Lee [13].

Unfortunately, theorem 1 does not provide any clue on the smallest dimension above which the diameter of associahedra is always twice the dimension minus four. The diameters of these polytopes have been obtained up to dimension 15 using a computer program, showing that this smallest dimension is not less than 9 (see page 654 in [17]). In addition, associahedra of dimension 10 to 15 were found to satisfy the rule stated by theorem 1, leading to the following conjecture, also originally formulated in [17] in the context of triangulations and flips: *the diameter of the d-dimensional associahedron is $2d-4$ when d is greater than* 9. The problem remained open until now, and made its way to textbooks [5,7].

While the statement of theorem 1 is purely combinatorial, the proof given in [17] is based on hyperbolic geometry. The authors of this paper comment that the role played by geometry in this problem “may seem mysterious” and they add that there should exist a proof using only combinatorial arguments. These observations led to consider the existence of a combinatorial proof for theorem 1 as an open problem on its own [4,7]. Recently, Patrick Dehornoy made progress towards such a proof by obtaining a lower bound of the form $2d-O(\sqrt{d})$ on the diameter of the d-dimensional associahedron, using purely combinatorial arguments [4].

The two open problems mentioned above are solved in this article. In other words, a combinatorial proof is given that the diameter of the d-dimensional associahedron is $2d-4$ when d is greater than 9.

This result will be obtained using flip operations in the triangulations of convex polygons. A combinatorial description of these notions will be given in section 2 using the finite subsets of $\mathbb{Z}$. While the proofs could also be written using traditional 2-dimensional polygons, this alternative formalism will simplify the exposition of the results throughout the article. The correspondence found by Carl Lee between the 1-skeletons of associahedra and the flip-graphs of polygons is also given in section 2 using this formalism.

Sections 3, 4, and 5 are devoted to stating general equalities and inequalities on the distance between two triangulations in the flip-graph of a polygon. The first of these inequalities, found in section 3 generalize lemma 3 from [17]. The inequalities given in section 4 are based on the operation of contracting an edge in a triangulation. It will be shown in particular that contracting an edge in the triangulations of a polygon induces contractions in the flip-graph of this polygon. Section 5 further provides equalities on the distances in two distinct flip-graphs, when the triangulations of a polygon are mapped into triangulations of another polygon in an adequate way.

Three pairs of triangulations of a polygon with n vertices are described in section 6 when n is greater than 2, along with some of their properties. A recursive lower bound on the flip distance of the first pair is found in section 7 using the results of sections 3 to 6. The two other pairs will serve as intermediates for the statement of the results. The diameters of associahedra are obtained in section 8 based on this recursive lower bound. In section 9, a third related open problem, posed in [4], is solved.

A discussion on the possibility of finding similar results on some generalizations of associahedra [9, 1, 2] completes the article.

2 Flip-graphs of polygons and 1-skeletons of associahedra

In this article, an *edge* on a set S is a subset of S with two elements. The elements of an edge will also be referred to as its *vertices.* The finite subsets of $\mathbb{Z}$ provide a combinatorial representation of convex polygons that allows for a convenient treatment of their triangulations. Indeed, if one identifies the vertex set of a convex polygon with a finite set $\pi \subset \mathbb{Z}$ in such a way that every two consecutive elements of π are the vertices of a boundary edge, the properties of edges on polygons can be stated combinatorially. For instance the notion of crossing edges can be defined as follows on $\mathbb{Z}$:

Definition 1 Two edges ε and ς on $\mathbb{Z}$ are *crossing* if they are disjoint and exactly one vertex of ε is strictly between the two vertices of ς.

Consider a finite subset π of $\mathbb{Z}$. The elements of π will be, as in the case of edges, called its vertices. An edge on π will be called a *boundary edge* on π if it does not cross any edge on π, and an *interior edge* on π otherwise. In addition, the set of all boundary edges on π will be denoted by $\beta(\pi)$. Following definition 1, the elements of $\beta(\pi)$ are the edges whose vertices are consecutive elements of π, and the edge whose vertices are the smallest and the greatest elements of π. Using the notion of crossing edges, one can directly transpose the usual definition of triangulations from 2-dimensional convex polygons to finite subsets of $\mathbb{Z}$.

Definition 2 Let π be a finite subset of $\mathbb{Z}$. A triangulation of π is a set of pairwise non-crossing edges on π that is maximal for the inclusion.

Because of the above-mentioned identification, any finite subset of $\mathbb{Z}$ with at least three elements is called a *polygon* hereafter. Observe that definitions 1 and 2 are not stated for polygons only, but also for the subset of $\mathbb{Z}$ with less than three elements. This will avoid unnecessary technicalities in the following sections. This alternative formalism for polygons and their triangulations not only provides a combinatorial representation of these objects, but also a natural ordering of their vertices, and the possibility to directly use these vertices in arithmetic operations. Yet, they still can be represented as their 2-dimensional counterpart. The reader is referred to [5] for a comprehensive study of triangulations in general.

Now consider a triangulation T of a polygon π. A classical (and straightforward) result states that all the boundary edges on π belong to T. Hence, the boundary edges on π will also be referred to as the boundary edges *of* T. Similarly, the interior edges on π that belong to T will be called the interior edges *of* T. A bijection α from a polygon π onto a polygon π' is called *boundary-preserving* if it induces a bijection from $\beta(\pi)$ onto $\beta(\pi')$. In other words, α is boundary-preserving if it commutes with β. Note in particular that any strictly monotonic map from a polygon onto another one is boundary-preserving. According to the following classical proposition, boundary-preserving maps maintain the whole structure of a triangulation:

Proposition 1 *A boundary-preserving bijection from a polygon π onto a polygon π' maps every triangulation of π to a triangulation of π'.*

This proposition follows from the fact that two edges on a polygon π are crossing if and only if their images by a boundary-preserving bijection from π onto any given polygon are crossing. The next proposition is a classical characterization of triangulations using their number of edges:

Proposition 2 *Let n be an integer greater than 1 and π a finite subset of $\mathbb{Z}$ with cardinality n. A set of pairwise non-crossing edges on π is a triangulation of π if and only if it has cardinality $2n-3$.*

As first observed by Carl Lee [13], flip-graphs of polygons are isomorphic to the 1-skeletons of associahedra. These graphs are now defined. Consider an edge ε on a polygon π, and a triangulation T of this polygon. The following subset of π is called the *link* of ε in T:

$$\lambda(\varepsilon, T) = \{x \in \pi : \forall\, y \in \varepsilon,\ \{x, y\} \in T\}.$$

According to well-known properties of triangulations [5], ε is a boundary edge of T if and only if $\lambda(\varepsilon, T)$ is a singleton. Note that this only holds when π is a polygon (otherwise $\lambda(\varepsilon, T)$ is necessarily empty). Now assume that ε is an interior edge of T. In this case, $\lambda(\varepsilon, T)$ is another interior edge on π that crosses ε. In other words, the set $\varepsilon \cup \lambda(\varepsilon, T)$ is a quadrilateral whose two diagonals are ε and its link in T. Consider the set $\mathfrak{F}(\varepsilon, T)$ obtained replacing ε by $\lambda(\varepsilon, T)$ in T:

$$\mathfrak{F}(\varepsilon, T) = [T \setminus \{\varepsilon\}] \cup \{\lambda(\varepsilon, T)\}.$$

Another straightforward result on triangulations is that $\mathfrak{F}(\varepsilon, T)$ is also a triangulation of π. The operation of replacing ε by $\lambda(\varepsilon, T)$ in T is called a *flip* [5]. In other words, when all the boundary edges of a quadrilateral q are edges of T, the flip operation consists in replacing the diagonal of q that belongs to T by the other diagonal of q. In particular the link of $\lambda(\varepsilon, T)$ in $\mathfrak{F}(\varepsilon, T)$ is ε itself and T can be recovered by flipping $\lambda(\varepsilon, T)$ in $\mathfrak{F}(\varepsilon, T)$. The graph whose vertices are the triangulations of a polygon π and whose edges connect two triangulations whenever they can be obtained from one another by a flip is called the *flip-graph* of π. Two triangulations of π will be called *adjacent* when they are the vertices of an edge in the flip-graph of π.

As mentioned above, the following theorem is proven in [13]:

Theorem 2 *Let d be a non-negative integer and π a polygon with $d+3$ vertices. The 1-skeleton of the d-dimensional associahedron is isomorphic to the flip-graph of π.*

According to this theorem, results on the flip-graphs of polygons can be transposed from and to the 1-skeletons of associahedra. In particular the flip-graph of a polygon is always connected (this connectedness property can be obtained from a much simpler argument, though, see [17]). In other words, one can obtain a triangulation of a polygon π by performing a sequence of flips from any other triangulation of π. Note that flip-graphs are not always connected in more general cases [5].

3 First equalities and inequalities on flip distances

Results on the flip-graphs of polygons, and especially regarding their diameters, can be found in [17]. These results will be needed later the paper and the beginning of this section is devoted to stating them. In the remainder of the section, one of these results is generalized. For any positive integer n, denote by $\iota(n)$ the set of the n smallest non-negative integers:

$$\iota(n) = \{0, 1, ..., n - 1\}.$$

Consider a finite subset π of $\mathbb{Z}$ and denote by $\tau(\pi)$ the set of all the triangulations of π. Further consider two triangulations U and V of π. A *path* of length k from U to V is a map $\psi : \iota(k + 1) \to \tau(\pi)$ so that $\psi(0) = U$, $\psi(k) = V$, and for all $i \in \iota(k)$, triangulations $\psi(i)$ and $\psi(i + 1)$ are adjacent. A path from U to V is called *minimal* when its length is minimal among all the paths from U to V. The *flip distance* of U and V, denoted by $\delta(\{U, V\})$ is the length of any minimal path between U and V. Lemma 2 in [17] states the following inequality on flip distances:

Lemma 1 *Let U and V be two triangulations of a polygon π with n vertices. If n is greater than 12, then $\delta(\{U, V\}) \leq 2n - 10$.*

This lemma is proven in [17] using purely combinatorial arguments. Invoking theorem 2, lemma 1 can be rephrased as: *the diameter of the d-dimensional associahedron is not greater than $2d - 4$ when d is greater than 9.* One can see that, if the conjecture on the diameter of associahedra is true, this bound is sharp. As discussed in [17, 4, 5], finding lower bounds on flip distances is the difficult part, tough. Lemma 3 from [17], whose proof is combinatorial as well, will be helpful in the search for lower bounds. Part (a) of this lemma can be translated as follows using the formalism introduced above:

Lemma 2 *Let U and V be two triangulations of a polygon π and ε an interior edge of U. If $\lambda(\varepsilon, U) \in V$, then $\delta(\{U, V\}) = \delta(\{\mathfrak{F}(\varepsilon, U), V\}) + 1$.*

Consider two triangulations U and V of a same polygon. Lemma 2 states that if an edge of V can be introduced in U by a flip, then there is a minimal path from U to V that begins with this flip. Part (b) of the same lemma found in [17] states that, when an edge has been removed along a minimal path, it cannot reappear later along the same path:

Lemma 3 *Let ψ be a minimal path of length k between two triangulations of a polygon π. If an edge ε on π is an edge of both $\psi(0)$ and $\psi(k)$, then for every integer i in $\iota(k+1)$, ε is an edge of $\psi(i)$.*

Two generalizations of lemma 2 are now stated. Consider two triangulations U and V of the same polygon and an interior edge ε of U. According to the following first generalization, if the link of ε in U is an edge of some triangulation along any minimal path between U and V, then ε is the first edge flipped in some minimal path between these two triangulations.

Lemma 4 *Let ψ be a minimal path of length k between two triangulations of a polygon π, and ε an interior edge of $\psi(0)$. If there exists $i \in \iota(k+1)$ so that $\lambda(\varepsilon, \psi(0))$ is an edge of $\psi(i)$, then $\delta(\{\mathfrak{F}(\varepsilon, \psi(0)), \psi(k)\}) = k-1$.*

Proof Assume that there exists an integer i in $\iota(k+1)$ so that the link of ε in $\psi(0)$ is an edge of $\psi(i)$. Denoting triangulation $\mathfrak{F}(\varepsilon, \psi(0))$ by T, one immediately obtains from lemma 2 that:

$$\delta(\{T, \psi(i)\}) = i-1. \tag{1}$$

In addition, according to the triangular inequality,

$$\delta(\{T, \psi(i)\}) + \delta(\{\psi(i), \psi(k)\}) \geq \delta(\{T, \psi(k)\}). \tag{2}$$

Since ψ is a minimal path, $\psi(i)$ and $\psi(k)$ have flip distance $k-i$. Hence, eliminating $\delta(\{T, \psi(i)\})$ between (1) and (2) yields:

$$k-1 \geq \delta(\{T, \psi(k)\}). \tag{3}$$

Another triangular inequality states that the right-hand side of inequality (3) is not less than its left-hand side, which completes the proof. □

This lemma can in turn be generalized to sequences of flips. In particular, the next theorem states that, under requirements similar to the conditions of lemma 4, there exists a path between two triangulations of a same polygon whose first several flips can be prescribed.

Theorem 3 *Let ψ be a minimal path of length k between two triangulations of a polygon π, ϕ a map from $\iota(j+1)$ into $\tau(\pi)$, $(\varepsilon_i)_{i\in\iota(j)}$ a sequence of interior edges on π, and $\alpha : \iota(j) \rightarrow \iota(k+1)$ an increasing map so that for all $i \in \iota(j)$, the following three assertions hold:*

i. ε_i *is an edge of* $\phi(i)$,
ii. $\phi(i+1) = \mathfrak{F}(\varepsilon_i, \phi(i))$,
iii. $\lambda(\varepsilon_i, \phi(i)) \in \psi \circ \alpha(i)$.

If $\phi(0) = \psi(0)$, then ϕ is a minimal path from $\phi(0)$ to $\phi(j)$, and the flip distance of $\phi(j)$ and $\psi(k)$ is equal to $k-j$.

Proof The theorem will be proven by induction on j. First observe that if $j=1$, then the statement of the theorem simplifies to that of lemma 4 when one respectively replaces ε_0, $\phi(0)$, and $\phi(1)$ by ε, $\psi(0)$, and $\mathfrak{F}(\varepsilon, \psi(0))$. Hence the result holds in this case.

Now assume that the result holds for a given positive integer j and for any value of k. Consider a minimal path ψ of length k between two triangulations of π, a map ϕ from $\iota(j+2)$ into $\tau(\pi)$, a sequence $(\varepsilon_i)_{i\in\iota(j+1)}$ of interior edges on π, and an increasing map α from $\iota(j+1)$ into $\iota(k+1)$. Assume in addition that for every integer i in $\iota(j+1)$, the three assertions in the statement of the theorem hold. In this case, the restriction of ψ to the smallest $\alpha(j)+1$ non-negative integers is a minimal path from $\psi(0)$ to $\psi\circ\alpha(j)$. Moreover, the restriction of α to $\iota(j)$ is an increasing map and the three statements in the lemma are still satisfied for every integer i in $\iota(j)$. Therefore, by induction, the following equality holds:

$$\delta(\{\phi(j), \psi\circ\alpha(j)\}) = \alpha(j) - j.$$

It is also obtained by induction that the restriction of ϕ to $\iota(j+1)$ is a minimal path from $\phi(0)$ to $\phi(j)$. As a consequence, one can find a path ψ' of length k from $\psi(0)$ to $\psi(k)$ that coincides with path ϕ on $\iota(j+1)$ and with path ψ on $\iota(k+1)\setminus\iota\circ\alpha(j)$.

As every path of length k between $\psi(0)$ and $\psi(k)$ is a minimal path, then ψ' is a minimal path between these two triangulations. Hence its restriction to $\iota(k+1)\setminus\iota(j)$ induces a minimal path of length $k-j$ from $\phi(j)$ to $\psi(k)$. As ψ' and ψ coincide in $\alpha(j)$, and as the link of ε_j in $\phi(j)$ belongs to $\psi\circ\alpha(j)$, it follows from lemma 4 that:

$$\delta(\{\mathfrak{F}(\varepsilon_j, \phi(j)), \psi(k)\}) = k - j - 1. \tag{4}$$

Invoking the second assertion with $i=j$, one obtains from equation (4) that the flip distance of triangulations $\phi(j+1)$ and $\psi(k)$ is $k-(j+1)$. As a consequence, there exists minimal paths from $\psi(0)$ to $\psi(k)$ whose restrictions to $\iota(j+2)$ coincide with ϕ. It immediately follows that ϕ is a minimal path from triangulation $\phi(0)$ to triangulation $\phi(j+1)$. □

4 Inequalities induced by contractions

Additional inequalities on flip distances are stated in this section, based on the operation of contracting an edge in the triangulations of a polygon π. Informally, this operation consists in bringing together consecutive vertices of π. Performing a contraction in every triangulation of π results in the triangulations of a smaller polygon π'. It will be shown that this operation induces a continuous surjection from the flip-graph of π onto that of π', and therefore makes it possible to compare the distances in these two flip-graphs. The inequalities thus obtained will be instrumental in sections 6 to 8.

4.1 Contracting an edge in the triangulations of a polygon

Consider an edge κ on $\mathbb{Z}$ and denote by a and b its vertices with the requirement that $a<b$. Let $\Lambda_\kappa:\mathbb{Z}\rightarrow\mathbb{Z}$ be the map so that for every relative integer x,

$$\Lambda_\kappa(x)=\begin{cases}\min(\{x,a\}) & \text{if } x\leq b,\\ x-(b-a) & \text{otherwise.}\end{cases}$$

Further denote by $\overline{\kappa}$ the intersection of $\mathbb{Z}$ with $[a,b]$. Intuitively, Λ_κ *contracts* $\overline{\kappa}$ to its minimum element, decreases by $b-a$ the integers greater than b, and leaves the integers less than a unaffected. Observe that Λ_κ is a monotonically increasing map on $\mathbb{Z}$. In addition, two distinct elements of $\mathbb{Z}$ have the same image by Λ_κ if and only if they both belong to $\overline{\kappa}$. In particular Λ_κ maps an edge ε on $\mathbb{Z}$ to an edge on $\Lambda_\kappa(\mathbb{Z})$ if and only if ε is not an edge on $\overline{\kappa}$. Let E be the set of all the edges on $\overline{\kappa}$.

It is used as a convention in the following that, if Λ_κ is defined on all the elements of a set S then $\Lambda_\kappa(S)$ is the set $\{\Lambda_\kappa(s):s\in S\}$. Now consider a triangulation T of a polygon π and denote:

$$\Lambda^\star_\kappa(T)=\Lambda_\kappa(T\setminus E),$$

As $\Lambda^\star_\kappa$ first removes the edges on $\overline{\kappa}$ from T, then its image of T only contains edges. Now recall that $\beta(\pi)$ is precisely made up of the edges whose vertices are consecutive elements of π, and of the edge ε whose vertices are the smallest and the largest elements of π. Denote $\beta^\star(\pi)=\beta(\pi)\setminus\{\varepsilon\}$ and assume that κ belongs to $\beta^\star(\pi)$. In this case, the link of κ in T contains a unique element c and the only two edges of $T\setminus E$ that have the same image by Λ_κ are $\{a,c\}$ and $\{b,c\}$ (recall that the two vertices of κ have been denoted by a and b). Hence $\Lambda^\star_\kappa(T)$ has exactly one element less than $T\setminus E$, and two less than T because κ is the unique edge on $\overline{\kappa}$ that belongs to T. It follows from these observations that for any edge κ on π, $\Lambda^\star_\kappa$ defines a map from the triangulations of π into the triangulations of $\Lambda_\kappa(\pi)$:

Proposition 3 *Let T be a triangulation of a polygon π. For any edge κ on π, the image of T by $\Lambda^\star_\kappa$ is a triangulation of $\Lambda_\kappa(\pi)$.*

Proof The result is first proven when κ belongs to $\beta^\star(\pi)$. In this case, κ is the unique edge on $\overline{\kappa}$ contained in T and the definition of $\Lambda^\star_\kappa(T)$ simplifies to:

$$\Lambda^\star_\kappa(T)=\Lambda_\kappa(T\setminus\{\kappa\}).$$

Consider two edges ε and ς in T and assume that they are both distinct from κ. It follows from this assumption that the images of these two edges by Λ_κ are edges on $\Lambda_\kappa(\pi)$. As T is a triangulation, then ε and ς are non-crossing. Hence, according to definition 1, these edges are are either non-disjoint, or each of them has an even number of its vertices (i.e. neither or both) between the vertices of the other edge. This property is preserved by map Λ_κ because this map is monotonically increasing. As a consequence, edges $\Lambda_\kappa(\varepsilon)$ and $\Lambda_\kappa(\varsigma)$ are non-crossing. Hence, $\Lambda^\star_\kappa(T)$ is a set of non-crossing edges on $\Lambda_\kappa(\pi)$.

Recall that the only two distinct elements of π with the same image by $\varLambda_\kappa$ are the vertices of κ. Therefore, if n denotes the cardinality of π, then $\varLambda_\kappa(\pi)$ has exactly $n-1$ elements. According to proposition 2, T contains exactly $2n-3$ edges. As $\varLambda_\kappa^\star(T)$ has two elements less than T, then it contains $2(n-1)-3$ edges, and it follows from the same proposition that $\varLambda_\kappa^\star(T)$ is a triangulation of $\varLambda_\kappa(\pi)$.

Now assume that κ is any edge on π. The result is obtained by induction on the cardinality of $\overline{\kappa}\cap\pi$. Observe that this cardinality is not less than 2 and that it is equal to 2 if and only if $\kappa\in\beta^\star(\pi)$. Hence the result has already been obtained when $\overline{\kappa}\cap\pi$ has 2 elements.

Assume that for some integer k greater than 1, the desired property holds whenever $\overline{\kappa}\cap\pi$ has cardinality k. Further assume that $\overline{\kappa}\cap\pi$ has $k+1$ vertices. Consider the edge ς whose vertices are the two greatest elements of $\overline{\kappa}\cap\pi$ and denote $\upsilon=\varLambda_\varsigma(\kappa)$. According to this construction, ς is an element of $\beta^\star(\pi)$ and $\overline{\upsilon}\cap\pi$ has cardinality k.

In addition, $\varLambda_\kappa$ and $\varLambda_\kappa^\star$ can be decomposed as follows:

$$\varLambda_\kappa=\varLambda_\upsilon\circ\varLambda_\varsigma \text{ and } \varLambda_\kappa^\star=\varLambda_\upsilon^\star\circ\varLambda_\varsigma^\star.$$

As ς belongs to $\beta^\star(\pi)$ then, as shown above, $\varLambda_\varsigma^\star(T)$ is a triangulation of $\varLambda_\varsigma(\pi)$. If $\varLambda_\varsigma(\pi)$ is not a polygon (i.e. if it has cardinality below 3), then $\varLambda_\kappa(\pi)$ is a singleton and $\varLambda_\kappa^\star(T)$ is empty. In this case, the desired result immediately follows from definition 2. Now, assume that $\varLambda_\varsigma(\pi)$ is a polygon. As $\overline{\upsilon}\cap\pi$ has cardinality k, it is obtained by induction that $\varLambda_\upsilon^\star\circ\varLambda_\varsigma^\star(T)$ is a triangulation of $\varLambda_\upsilon\circ\varLambda_\varsigma(\pi)$. Since maps $\varLambda_\upsilon\circ\varLambda_\varsigma$ and $\varLambda_\upsilon^\star\circ\varLambda_\varsigma^\star$ respectively coincide with $\varLambda_\kappa$ and $\varLambda_\kappa^\star$, this completes the proof. □

4.2 Contractions induced in the flip-graph of a polygon

The way flip-graphs and the paths within them react to contractions is now investigated. Consider a path ψ of length k between two triangulations of a polygon π. By definition, there exists a sequence $(\varepsilon_i)_{i\in\iota(k)}$ of interior edges on π so that for every integer i in $\iota(k)$, $\varepsilon_i\in\psi(i)$ and $\psi(i+1)=\mathfrak{F}(\varepsilon_i,\psi(i))$. For any edge κ on π, denote by $\gamma_\kappa(\psi)$ the set of all the elements i of $\iota(k)$ so that quadrilateral $\varepsilon_i\cup\lambda(\varepsilon_i,\psi(i))$ contains at least two vertices of $\overline{\kappa}$:

$$\gamma_\kappa(\psi)=\{i\in\iota(k):|\overline{\kappa}\cap[\varepsilon_i\cup\lambda(\varepsilon_i,\psi(i))]|\geq 2\}.$$

Observe that flipping ε_i in $\psi(i)$ modifies the links of all the edges on quadrilateral $\varepsilon_i\cup\lambda(\varepsilon_i,\psi(i))$ and leaves the other links unaffected. Hence, an integer i belongs to $\gamma_\kappa(\psi)$ if and only if the flip that transforms $\psi(i)$ into $\psi(i+1)$ modifies the link in $\psi(i)$ of at least one edge on $\overline{\kappa}$. It will be shown (theorem 4 below) that the cardinality of $\gamma_\kappa(\psi)$ is the number of flips that disappear from ψ when contracting κ in all the triangulations visited by ψ. The following lemma provides a first result on the cardinality of this set:

Lemma 5 *Let ψ be a minimal path between two triangulations U and V of a polygon π. An element κ of $\beta^\star(\pi)$ has distinct links in U and in V if and only if $\gamma_\kappa(\psi)$ is non-empty.*

Proof Let κ be an element of $\beta^\star(\pi)$. Denote by a and b the two vertices of κ and by c the unique element of its link in U. Further denote by k the length of ψ, and for every $i \in \iota(k)$, call ε_i the interior edge of $\psi(i)$ so that:

$$\psi(i+1) = \mathfrak{F}(\varepsilon_i, \psi(i)).$$

Consider the set S of the integers i in $\iota(k)$ so that ε_i is equal to either $\{a, c\}$ or $\{b, c\}$. Observe that if $\gamma_\kappa(\psi)$ is non-empty then its smallest element necessarily belongs to S. Inversely, if S is non-empty, then its smallest element belongs to $\gamma_\kappa(\psi)$. In other words, a flip along path ψ removes edge $\{a, c\}$ or edge $\{b, c\}$ if and only if $\gamma_\kappa(\psi)$ is non-empty.

Assume that the link of κ in V is $\{c\}$. In this case, edges $\{a, c\}$ and $\{b, c\}$ both belong to U and to V. As ψ is a minimal path from U to V, it follows from lemma 3 that these two edges both belong to $\psi(i)$ for all $i \in \iota(k+1)$. Hence, these edges are not removed by any flip along path ψ, showing that $\gamma_\kappa(\psi)$ is empty. This provides an indirect proof of the first implication.

Now assume that the link of κ in V is distinct from $\{c\}$. In this case, $\{a, c\}$ and $\{b, c\}$ cannot both belong to V. Hence, at least one of these edges is removed by a flip along path ψ. As a consequence $\gamma_\kappa(\psi)$ is non-empty, which proves the second implication. □

Consider an edge κ on a polygon π and a path ψ between two triangulations U and V of this polygon. According to the next theorem, there exists a path ϕ between $\Lambda_\kappa(U)$ and $\Lambda_\kappa(V)$ whose length is obtained subtracting the cardinality of $\gamma_\kappa(\psi)$ from the length of ψ. In fact, path ϕ will be found by contracting edge κ in all the triangulations visited by path ψ and by subsequently removing unnecessary triangulations from the sequence thus obtained.

Theorem 4 *Let ψ be a path of length k between two triangulations U and V of a polygon π. For any edge κ on π, there exists a path of length $k - |\gamma_\kappa(\psi)|$ between triangulations $\Lambda^\star_\kappa(U)$ and $\Lambda^\star_\kappa(V)$.*

Proof Consider an edge κ on π. For all i in $\iota(k)$ denote by ε_i and ς_i the two edges so that $\psi(i+1)$ is obtained replacing ε_i by ς_i in $\psi(i)$. Further consider an arbitrary element j of $\iota(k)$ and denote quadrilateral $\varepsilon_j \cup \varsigma_j$ by q. It will be shown that, if j belongs to $\gamma_\kappa(\psi)$, then triangulations $\psi(j)$ and $\psi(j+1)$ are identical, and that otherwise, these triangulations are adjacent.

First assume that $j \in \gamma_\kappa(\psi)$. It follows that at least two vertices of quadrilateral q belong to $\overline{\kappa}$. In particular, $\Lambda_\kappa(q)$ has 1, 2, or 3 elements. As in addition, $\overline{\kappa}$ is made up of consecutive integers, then $\overline{\kappa}$ admits at least one boundary edge on q as a subset. In this case, $\Lambda_\kappa(\varepsilon_i)$ and $\Lambda_\kappa(\varsigma_i)$ can each alternatively be obtained as the image by Λ_κ of a boundary edge on quadrilateral q. As triangulations $\psi(j)$ and $\psi(j+1)$ both contain all the boundary edges of q and



Fig. 1 The flip-graph of the hexagon (solid lines) and the equivalence classes of $\sim_\kappa$ (dotted lines), where κ is the boundary edge at the top of the hexagon.

as they are obtained from one another by exchanging ε_j and ς_j, then these triangulations have the same image by $\Lambda_\kappa^\star$.

Now assume that $j \notin \gamma_\kappa(\psi)$. In this case, quadrilateral q has at most one vertex in $\overline{\kappa}$. Hence, $\Lambda_\kappa(q)$ is still a quadrilateral, and the restriction of Λ_κ to q is strictly increasing. It follows that the images of ε_i and ς_i by Λ_κ are the diagonals of $\Lambda_\kappa(q)$. For the same reason, Λ_κ maps the boundary edges of q onto the boundary edges of $\Lambda_\kappa(q)$. As $\psi(j)$ contains all the boundary edges of q, this shows that $\Lambda_\kappa(\varsigma_j)$ is the link of $\Lambda_\kappa(\varepsilon_j)$ in $\Lambda_\kappa^\star \circ \psi(j)$. Now observe that $\Lambda_\kappa^\star \circ \psi(j+1)$ is obtained from $\Lambda_\kappa^\star \circ \psi(j)$ by replacing $\Lambda_\kappa(\varepsilon_j)$ with $\Lambda_\kappa(\varsigma_j)$. Therefore, these two triangulations are adjacent.

Hence, one can build a path ϕ from $\Lambda_\kappa^\star(U)$ to $\Lambda_\kappa^\star(V)$ that successively visits triangulations $\Lambda_\kappa^\star(\psi(i))$, where i ranges from 0 to k, skipping the elements of $\gamma_\kappa(\psi)$. As a consequence, path ϕ has length $k - |\gamma_\kappa(\psi)|$. □

Theorem 4 can be considered as the most important result of the section because it relates the flip distance of two given triangulations with that of two other triangulations with less vertices. A more general result on flip-graphs can be obtained from the main arguments in the proof of this theorem. Indeed, consider an edge κ on a polygon π and denote $U \sim_\kappa V$ whenever two triangulations U and V of π have the same image under $\Lambda_\kappa^\star$. Observe that this defines an equivalence relation $\sim_\kappa$ on $\tau(\pi)$. When κ is contracted, any two triangulations that are equivalent under this relation are changed into their common image by $\Lambda_\kappa^\star$. Moreover, as shown in the proof of theorem 4, this contraction preserves the adjacency of any two triangulations that are not equivalent under $\sim_\kappa$. It immediately follows that *the flip-graph of* $\Lambda_\kappa(\pi)$ *is isomorphic to the graph obtained contracting every element of* $\tau(\pi)/{\sim_\kappa}$ *in the flip-graph of* π. This property is illustrated in figure 1 in the case of the hexagon. In this figure the flip-graph of this polygon is represented using solid lines and its fourteen triangulations are explicitly depicted as the vertices of this graph. The dotted lines show the five elements of $\tau(\pi)/{\sim_\kappa}$, where κ is the boundary edge at the top of the hexagon. As can be seen, it is implicitly assumed that κ is not the edge whose vertices are the smallest and the greatest elements of the hexagon, otherwise $\sim_\kappa$ would admit a unique equivalence class. One can see that contracting the elements of $\tau(\pi)/\!\sim_\kappa$ in the flip-graph of the hexagon results in a simple cycle whose vertices are the five triangulations of the pentagon and whose edges correspond to the flips between these triangulations. While this property will not be explicitly needed in the following, it provides intuition on the meaning of theorem 4.

4.3 Further inequalities on flip distances

For any two triangulations U and V of a polygon π, denote by $\vartheta_\kappa(\{U,V\})$ the maximal cardinality of $\gamma_\kappa(\psi)$ over all the minimal paths ψ between U and V. The following inequality on flip distances is a consequence of theorem 4:

Corollary 1 *For any pair P of triangulations of a polygon π and any edge κ on π, $\delta(P) \geq \delta \circ \Lambda_\kappa^\star(P) + \vartheta_\kappa(P)$.*

Proof Consider a pair P of triangulations of a polygon π and an edge κ on π. By definition, there exists a minimal path ψ between the two triangulations in P so that $\gamma_\kappa(\psi)$ has cardinality $\vartheta_\kappa(P)$. It then follows from theorem 4 that there exists a path of length $\delta(P) - \vartheta_\kappa(P)$ between the two triangulations in pair $\Lambda_\kappa^\star(P)$. Hence, minimal paths between these two triangulations have length at most $\delta(P) - \vartheta_\kappa(P)$, which completes the proof. ☐

Consider a pair P of triangulations of a polygon π. One can see that corollary 1 provide better lower bounds on $\delta(P)$ when map ϑ_κ takes large values for some edge κ on π. If, however, this map only takes small values, then one can obtain alternative arguments as, for instance, the following theorem:

Theorem 5 *Let U and V be two triangulations of a polygon π, ψ a minimal path of length k from U to V, and κ an element of $\beta^\star(\pi)$. If $\gamma_\kappa(\psi)$ admits a unique element j, then the following three assertions hold:*

i. *For all $i \in \iota(j+1)$, $\lambda(\kappa, \psi(i))$ is equal to $\lambda(\kappa, U)$,*
ii. *For all $i \in \iota(k+1) \setminus \iota(j+1)$, $\lambda(\kappa, \psi(i)) = \lambda(\kappa, V)$,*
iii. *$\lambda(\kappa, U) \cup \lambda(\kappa, V)$ belongs to $\psi(j)$ and to $\psi(j+1)$.*

Proof Assume that j is the unique element of $\gamma_\kappa(\psi)$. Denote by a and b the two vertices of κ. As κ is a boundary edge on π, then $\lambda(\kappa, U)$ and $\lambda(\kappa, U)$ each contain exactly one element, respectively denoted by c and d. As $\gamma_\kappa(\psi)$ is non-empty, lemma 5 states that $c \neq d$.

Since ψ is a minimal path from U to V, then its restriction ϕ to $\iota(j+1)$ is a minimal path from U to $\psi(j)$ and its restriction to $\iota(k+1) \setminus \iota(j+1)$ induces a minimal ϕ' from $\psi(j+1)$ to V. As j is the only element of $\gamma_\kappa(\psi)$, then $\gamma_\kappa(\phi)$ and $\gamma_\kappa(\phi')$ are both empty. It then follows from lemma 5 that the links of ε in $\psi(j)$ and in $\psi(j+1)$ are respectively $\{c\}$ and $\{d\}$. Hence edges $\{a, c\}$ and $\{b, c\}$ both belong to triangulations U and $\psi(j)$. As ϕ is a minimal path between these triangulations, it follows from lemma 3 that $\{a,c\}$ and $\{b,c\}$ are edges of $\psi(i)$ for all $i \in \iota(j+1)$. As a consequence, for any such value of i, the link of ε in $\psi(i)$ is $\{c\}$, and the first assertion is proven.

Similarly, edges $\{a,d\}$ and $\{b,d\}$ both belong to triangulations $\psi(j+1)$ and V. As ϕ' is a minimal path between these triangulations, one obtains from lemma 3 that for all $i \in \iota(k+1) \setminus \iota(j+1)$, triangulation $\psi(i)$ contains $\{a,d\}$ and $\{b,d\}$. Hence, the link of ε in $\psi(i)$ is $\{d\}$ for all $i \in \iota(k+1) \setminus \iota(j+1)$, and the second assertion follows.

It has been shown that the links of $\{a,b\}$ in $\psi(j)$ and in $\psi(j+1)$ are respectively $\{c\}$ and $\{d\}$. As $c \neq d$, the flip that transforms $\psi(j)$ onto $\psi(j+1)$ exchanges the diagonals of quadrilateral $\{a,b,c,d\}$. Therefore, all the boundary edges of this quadrilateral belong to $\psi(j)$ and to $\psi(j+1)$. In particular $\{c,d\}$ is an edge of these two triangulations, which proves the third assertion. □

Consider a polygon π, an element κ of $\beta^\star(\pi)$, and a minimal path ψ between two triangulations of π. Theorem 5 states that, if $\gamma_\kappa(\psi)$ is a singleton, then each triangulations visited along path ψ contains given edges. Because of the placement of these edges, and since two edges of a triangulation are necessarily non-crossing, one can invoke theorem 5 to obtain lower bounds on map ϑ_κ:

Corollary 2 *Let U and V be two triangulations of a polygon π. If κ and υ are two elements of $\beta^\star(\pi)$ so that there exist $x \in \kappa$ and $y \in \upsilon$ satisfying:*

i. $\{x\} \cup \lambda(\kappa, U)$ *crosses* $[\upsilon \setminus \{y\}] \cup \lambda(\upsilon, V)$,
ii. $\{y\} \cup \lambda(\upsilon, U)$ *crosses* $[\kappa \setminus \{x\}] \cup \lambda(\kappa, V)$,
iii. $\lambda(\upsilon, U)$ *is not a subset of* $\{x\} \cup \lambda(\kappa, U)$,

then the images of $\{U, V\}$ by maps ϑ_κ and ϑ_υ are not both less than 2.

Proof Consider two edges κ and υ in $\beta^\star(\pi)$, and two vertices x and y of these respective edges that satisfy the assertions in the statement of the corollary. It follows from the first two assertions that κ has distinct links in U and in V. Otherwise, $\{x\} \cup \lambda(\kappa, U)$ and $[\upsilon \setminus \{y\}] \cup \lambda(\upsilon, V)$ are two crossing edges contained in V. For the same reason, υ has distinct links in U and V.

Consider a minimal path ψ from U to V. As edges κ and υ both have distinct links in triangulations U and V, it follows from lemma 5 that $\gamma_\kappa(\psi)$ and $\gamma_\upsilon(\psi)$ are non-empty. It will be shown indirectly that one of these two sets admits more than one element. Assume that $\gamma_\kappa(\psi)$ contains a unique element j and that $\gamma_\upsilon(\psi)$ contains a unique element l. According to theorem 5, the link of κ in $\psi(i)$ is equal to $\lambda(\kappa, U)$ for all $i \in \iota(j+1)$. Hence, edge $\{x\} \cup \lambda(\kappa, U)$ belongs to $\psi(i)$ for all $i \in \iota(j+1)$. One obtains from similar arguments that, $\{y\} \cup \lambda(\upsilon, U)$ is an edge of $\psi(i)$ for all $i \in \iota(l+1)$. Theorem 5 further states that $\psi(j+1)$ and $\psi(l+1)$ respectively contain edges $[\kappa \setminus \{x\}] \cup \lambda(\kappa, V)$ and $[\upsilon \setminus \{y\}] \cup \lambda(\upsilon, V)$. As the edges of a triangulation are non-crossing, it follows from the first assertion that $j \leq l$ and from the second one that $l \leq j$, which proves that j and l are identical.

It has been shown in the last paragraph that edges $\{x\} \cup \lambda(\kappa, U)$ and $\{y\} \cup \lambda(\upsilon, U)$ both belong to $\psi(j)$ and that edges $[\kappa \setminus \{x\}] \cup \lambda(\kappa, V)$ and

$[\upsilon \setminus \{y\}] \cup \lambda(\upsilon, V)$ both belong to $\psi(j+1)$. Hence, according to the first two assertions in the statement of the corollary, $\{x\} \cup \lambda(\kappa, U)$ and $\{y\} \cup \lambda(\upsilon, U)$ are edges of $\psi(j)$ that each cross an edge of $\psi(j+1)$. As $\psi(j)$ and $\psi(j+1)$ are adjacent, this shows that:

$$\{x\} \cup \lambda(\kappa, U) = \{y\} \cup \lambda(\upsilon, U).$$

This equality contradicts the third assertion in the statement of the corollary, providing an indirect proof that at least one of the two sets $\gamma_\kappa(\psi)$ and $\gamma_\upsilon(\psi)$ has more than one element. It immediately follows that the images of pair $\{U, V\}$ by ϑ_κ and ϑ_υ cannot both be less than 2. □

Under stronger conditions on triangulations U and V, the statement of corollary 2 can be simplified. The following weaker form of this corollary will actually be invoked most frequently in the remainder of this article:

Corollary 3 *Let U and V be two triangulations of a polygon π, and a, b, and c three vertices of this polygon. If the following asssertions hold:*

i. *$\{a, b\}$ and $\{b, c\}$ belong to $\beta^\star(\pi)$, and $\{a, c\}$ is an edge of V,*
ii. *$\lambda(\{a, b\}, U)$ and $\lambda(\{b, c\}, U)$ are distinct subsets of $\pi \setminus \{a, c\}$,*

then the images of $\{U, V\}$ by $\vartheta_{\{a,b\}}$ and $\vartheta_{\{b,c\}}$ are not both less than 2.

Proof Assume that U and V are two triangulations of a polygon π, and that a, b, and c are three vertices of π satisfying the two assertions in the statement of the corollary. Since $\{a, b\}$ and $\{b, c\}$ both belong to $\beta^\star(\pi)$, any edge on π that admits b as a vertex and whose other vertex belongs to $\pi \setminus \{a, c\}$ necessarily crosses $\{a, c\}$. In particular, edges $\{b\} \cup \lambda(\{a, b\}, U)$ and $\{b\} \cup \lambda(\{b, c\}, U)$ both cross edge $\{a, c\}$. In addition, as $\{a, c\}$ belongs to V then the links of edges $\{a, b\}$ and $\{b, c\}$ in this triangulation are respectively $\{c\}$ and $\{a\}$. Hence, calling $\kappa = \{a, b\}$, $\upsilon = \{b, c\}$, $x = b$, and $y = b$, the first two assertions in the statement of corollary 2 hold.

By hypothesis, $\lambda(\{a, b\}, U)$ and $\lambda(\{b, c\}, U)$ are distinct. As these two links are singletons, they are therefore disjoint. Moreover, $\lambda(\{b, c\}, U)$ is also disjoint from $\{b\}$. Hence, according to the notations chosen above, $\lambda(\upsilon, U)$ cannot be a subset of $\{x\} \cup \lambda(\kappa, U)$. In other words, the third assertion in the statement of corollary 2 also holds. It then follows from this corollary that $\vartheta_{\{a,b\}}(\{U, V\})$ and $\vartheta_{\{b,c\}}(\{U, V\})$ are not both less than 2. □

A triangle $\{a, b, c\}$ satisfying the first assertion in the statement of corollary 3 is sometimes called an *ear* of triangulation V. In fact, the usual definition of ears is slightly less restrictive. It is well known that any triangulation of a polygon admits at least one ear. The following proposition is closely related to this existence property. Its straightforward proof is provided nonetheless because of the difference between the first assertion in the statement of corollary 3 and the usual definition of ears:

Proposition 4 *Let π be a polygon and T a triangulation of π. There exist three vertices a, b, and c of π so that edges $\{a,b\}$ and $\{b,c\}$ belong to $\beta^\star(\pi)$ and edge $\{a,c\}$ belongs to T.*

Proof The proposition is proven by induction on $|\pi|$. If π has cardinality 3, then the result immediately follows by calling a, b, and c the vertices of π with the requirement that $a<b<c$.

Now assume that the result holds whenever π has cardinality at most k, where k is a given integer greater than 2. Further assume that π has $k+1$ elements. In this case, T admits at least one interior edge. Let κ be an interior edge of T. As κ is not a boundary edge of T then $3 \leq |\overline{\kappa}\cap\pi| \leq k$. In particular $\overline{\kappa}\cap\pi$ is a polygon. Since κ belongs to T, the edges on $\overline{\kappa}\cap\pi$ that belong to T form a triangulation U of $\overline{\kappa}\cap\pi$. Hence, by induction, there exist three vertices a, b, and c of $\overline{\kappa}\cap\pi$ so that edges $\{a,b\}$ and $\{b,c\}$ belong to $\beta^\star(\overline{\kappa}\cap\pi)$ and edge $\{a,c\}$ belongs to U. As $\beta^\star(\overline{\kappa}\cap\pi)$ and U are respectively a subset of $\beta^\star(\pi)$ and a subset of T, the result follows. □

This proposition will be used in the proof of the next theorem, that states additional lower bounds on the flip distance of two triangulations with given properties. These bounds are obtained by carrying out a well-chosen sequence of contractions in these triangulations, and by invoking corollary 3 between each of these contractions. Recall that $\beta^\star$ was exclusively defined on polygons. In order to avoid unnecessary technicalities when stating the following theorem, $\beta^\star(\varepsilon)$ has to be defined when ε is an edge on $\mathbb{Z}$ as the set $\{\varepsilon\}$.

Theorem 6 *Let U and V be two triangulations of a polygon π. If κ is an edge of V so that:*

i. For all $\varepsilon \in \beta^\star(\overline{\kappa}\cap\pi)$, $\lambda(\varepsilon,U)$ is disjoint from $\overline{\kappa}\cup\lambda(\kappa,V)$,
ii. Distinct elements of $\beta^\star(\overline{\kappa}\cap\pi)$ have distinct links in U,

then $\delta(\{U,V\}) \geq \delta\circ\varLambda^\star_\kappa(\{U,V\})+2|\overline{\kappa}\cap\pi|-3$.

Proof Consider an edge κ of V that satisfies the two assertions in the statement of the theorem. The result will be obtained by induction on the cardinality of $\overline{\kappa}\cap\pi$. Observe first that, if $\overline{\kappa}\cap\pi$ has exactly 2 elements, then κ is the unique element of $\beta^\star(\overline{\kappa}\cap\pi)$. Hence, according to the first assertion, its links in triangulations U and V are disjoint. In this case, it follows from lemma 5 that $\vartheta_\kappa(\{U,V\})$ is positive and, in turn from lemma 1 that:

$$\delta(\{U,V\}) \geq \delta\circ\varLambda^\star_\kappa(\{U,V\})+1.$$

As $2|\overline{\kappa}\cap\pi|-3$ is precisely equal to 1 in this case, then the result follows. Now assume that k is an integer greater than 1 so that the result holds if $\overline{\kappa}\cap\pi$ has cardinality k. Further assume that $\overline{\kappa}\cap\pi$ contains $k+1$ elements. As κ is an edge of V, then some triangulation T of $\overline{\kappa}\cap\pi$ is found as a subset of V. Observe that $\overline{\kappa}\cap\pi$ has at least three vertices. Hence it is a polygon and, according to proposition 4, it contains three vertices a, b, and c so that edges $\{a,b\}$ and $\{b,c\}$ belong to $\beta^\star(\overline{\kappa}\cap\pi)$ and edge $\{a,c\}$ belongs to T.

According to the second assertion, $\{a,b\}$ and $\{b,c\}$ have distinct links in U, and according to the first one, these links are subsets of $\pi \setminus \{a,c\}$. Hence, it follows from corollary 3 that the images of $\{U,V\}$ by $\vartheta_{\{a,b\}}$ and $\vartheta_{\{b,c\}}$ are not both less than 2. Denote by ς edge $\{a,b\}$ or edge $\{b,c\}$ with the requirement that $\vartheta_{\varsigma}(\{U,V\})$ is not less than 2. In this case, corollary 1 yields:

$$\delta(\{U,V\}) \geq \delta \circ \Lambda^{\star}_{\varsigma}(\{U,V\}) + 2. \tag{5}$$

Now observe that κ and ς are distinct. Indeed, b is a vertex of ς that does not belong to κ because it is strictly between two vertices of $\overline{\kappa}$ (i.e. a and c). In addition, $\Lambda^{\star}_{\varsigma}(V) = \Lambda_{\varsigma}(V \setminus \{\varsigma\})$ because ς is the only edge on π that belongs to V. Hence, the image of κ by Λ_{ς} is an edge of $\Lambda^{\star}_{\varsigma}(V)$. Denote by υ this edge. As ς is an element of $\beta^{\star}(\overline{\kappa} \cap \pi)$, then map Λ_{ς} induces a bijection $\beta^{\star}(\overline{\kappa}\cap\pi)\setminus\{\varsigma\} \rightarrow \beta^{\star}(\overline{\upsilon}\cap\pi)$ and another bijection $\pi\setminus\overline{\kappa} \rightarrow \Lambda_{\varsigma}(\pi)\setminus\overline{\upsilon}$. In addition, Λ_{ς} precisely maps $\overline{\kappa}$ to $\overline{\upsilon}$ and $\lambda(\kappa, V) \setminus \overline{\kappa}$ to $\lambda(\upsilon, \Lambda^{\star}_{\varsigma}(V)) \setminus \overline{\upsilon}$. Hence, the two assertions in the statement of the theorem still hold when one replaces κ, U, and V by respectively υ, $\Lambda^{\star}_{\varsigma}(U)$, and $\Lambda^{\star}_{\varsigma}(V)$. As ς belongs to $\beta^{\star}(\overline{\kappa} \cap \pi)$, then $\overline{\upsilon} \cap \pi$ has cardinality $|\overline{\kappa} \cap \pi| - 1$, and by induction:

$$\delta \circ \Lambda^{\star}_{\varsigma}(\{U,V\}) \geq \delta \circ \Lambda^{\star}_{\upsilon} \circ \Lambda^{\star}_{\varsigma}(\{U,V\}) + 2|\overline{\kappa} \cap \pi| - 5. \tag{6}$$

Finally observe that, by construction, $\Lambda^{\star}_{\upsilon} \circ \Lambda^{\star}_{\varsigma}$ coincides with $\Lambda^{\star}_{\kappa}$. As a consequence, combining inequalities (5) and (6) completes the proof. □

5 Injecting a triangulation into another one

It is shown in this section that transporting a pair of triangulation from a polygon to another one in an adequate manner preserves flip distance. Recall that, following proposition 1, the image of a triangulation by a boundary-preserving map remains a triangulation. As a direct consequence, composing a path with a boundary-preserving map results in a path:

Proposition 5 *Let α be a boundary-preserving bijection from a polygon π onto another polygon. If ψ is a path between two triangulations U and V of π then $\alpha \circ \psi$ is a path between $\alpha(U)$ and $\alpha(V)$.*

Proof Let ψ a minimal path of length k between two triangulations U and V of π. Consider an integer i in $\iota(k)$. According to proposition 1, $\alpha \circ \psi(i)$ and $\alpha \circ \psi(i+1)$ are two triangulations of $\alpha(\pi)$. Call ε and ς the two edges on π so that $\psi(i+1)$ is obtained replacing ε in $\psi(i)$ by ς. It follows that $\alpha \circ \psi(i+1)$ is obtained replacing $\alpha(\varepsilon)$ in $\alpha \circ \psi(i)$ by $\alpha(\varsigma)$. As α is boundary-preserving, the images of ε and ς by α are crossing edges, which proves that $\alpha \circ \psi(i)$ and $\alpha \circ \psi(i+1)$ are adjacent. As a consequence, $\alpha \circ \psi$ is a path between triangulations $\alpha(U)$ and $\alpha(V)$. □

It follows from this proposition that two triangulations of a given polygon π have the same flip distance as their images by a boundary-preserving bijection from π onto another polygon. One can generalize this result by considering maps that inject triangulations of π into triangulations of another polygon, instead of just boundary-preserving bijections. The following lemma states that, not only flip distances are preserved by such maps, but also the values of ϑ_κ for any edge κ in $\beta^\star(\pi)$:

Lemma 6 *Let U and V be two triangulations of a polygon π and α a boundary-preserving bijection from π onto a polygon $\alpha(\pi)$. If U' and V' are two triangulations of a polygon π' satisfying:*

i. *$\alpha(U)$ is a subset of U',*
ii. *$\alpha(V)$ is a subset of V',*
iii. *$U' \setminus \alpha(U) = V' \setminus \alpha(V)$,*

then pairs $\{U, V\}$ and $\{U', V'\}$ have the same flip distance, and for any edge κ in $\beta^\star(\pi)$, $\vartheta_\kappa(\{U, V\})$ is equal to $\vartheta_{\alpha(\kappa)}(\{U', V'\})$.

Proof Assume that U' and V' are two triangulations of a polygon π' so that the three assertions in the statement of the lemma hold. As α is a boundary-preserving bijection from π onto $\alpha(\pi)$ then, according to proposition 4, the map obtained composing a path between U and V with α is a path between triangulations $\alpha(U)$ and $\alpha(V)$.

As $\alpha(U)$ is a triangulation of $\alpha(\pi)$, it further follows from the first statement that removing every edge of $\alpha(U)$ from U' and replacing them by the edges of another triangulation of $\alpha(\pi)$ results in a triangulation of π'. Hence, according to the third statement, if ψ is a path of length k from U to V, then one can build a path $\tilde{\psi}$ of length k from U' to V' by taking the union of the triangulations along path $\alpha \circ \psi$ with set $U' \setminus \alpha(U)$.

In fact every minimal path from U' to V' is equal to $\tilde{\psi}$ for some path ψ from U to V. Indeed, according to the first two statements, U' and V' both admit a triangulation of $\alpha(\pi)$ as a subset. Hence, U' and V' both contain all the boundary edges of $\alpha(\pi)$. If further follows from the third statement that U' and V' both admit $U' \setminus \alpha(U)$ as a subset. Lemma 3 states that, in this case, $\beta \circ \alpha(\pi)$ and $U' \setminus \alpha(U)$ are subsets of every triangulation visited along any minimal path from U' to V'. As a consequence, every flip in such a path replaces an interior edge on $\alpha(\pi)$ by another interior edge on $\alpha(\pi)$. In particular, every minimal path from U' to V' is necessarily equal to $\tilde{\psi}$ for some path ψ from U to V.

Now observe that ψ and $\tilde{\psi}$ have the same length. Therefore, one obtains that $\tilde{\psi}$ is a minimal path from U' to V' if and only if ψ is a minimal path from U to V. This shows that $\delta(\{U, V\})$ is equal to $\delta(\{U', V'\})$. Finally, consider an edge κ in $\beta^\star(\pi)$. By construction, $\gamma_\kappa(\psi)$ and $\gamma_{\alpha(\kappa)}(\tilde{\psi})$ are identical. As a direct consequence, $\vartheta_\kappa(\{U, V\})$ and $\vartheta_{\alpha(\kappa)}(\{U', V'\})$ are also equal. □

6 Three pairs of triangulations

In this section, three pairs of triangulations of polygon $\iota(n)$ are defined. The first of these three pairs will be shown in the following sections to have flip distance $2n-10$ when n is greater than 12. The two other pairs will be used in the statement of intermediate results. The triangulations contained in the first pair are described informally in the following paragraph in order to provide some preliminary intuition.

Consider the triangulations of the hexagon shown in figure 1. The triangulation in the top left of the figure is made up of three interior edges that share a common vertex. A triangulation of this kind will be called a *comb*. More generally, when a vertex v of a triangulation T is incident to $k \geq 3$ interior edges of T, it will be said that T admits a comb with k *teeth* at vertex v. Now consider the the triangulation in the top right of figure 1. The interior edges of this triangulation form a simple path whose successive vertices alternate between two sides of the hexagon. Such a triangulation is called a *zigzag*. As was the case with combs, zigzags can contain an arbitrary large number of vertices (see for instance figure 2), and be found as a part of some triangulation. Note that zigzags and combs were already used in the context of binary trees in [4] to investigate the same problem. The two triangulations that will be shown to have maximal distance will be made up of a zigzag with small combs attached at both ends (see figure 3). These triangulations are not unrelated with the binary trees conjectured in [4] to have maximal rotation distance. This conjecture will be disproved in the last section, though.

In the following of the section, zigzags are first formally defined. Combs are then introduced at each end of these zigzags by means of two contractions, in order to build a pair of triangulations that will be proven maximally-distant in the next sections. Two other pairs of triangulations are subsequently defined, also by carrying out adequate contractions in zigzags. Several inequalities involving the flip distances of these three pairs are given throughout the section. These inequalities will be needed later in the article.

6.1 Zigzags

Let k by an arbitrary relative integer, and n an integer greater than 2. Denote by $f_n(k)$ the following set of edges on $\iota(n)$, where $\equiv$ refers to the usual congruence relation in the integers:

$$f_n(k) = \{\{i,j\} \subset \iota(n) : i \neq j,\, i+j \equiv k \pmod{n}\}.$$

Further consider the following two sets of edges on $\iota(n)$:

$$\left\{\begin{array}{l} Z_n^-(k) = f_n(2k-1) \cup f_n(2k) \cup \beta(\iota(n)), \\ Z_n^+(k) = f_n(2k) \cup f_n(2k+1) \cup \beta(\iota(n)). \end{array}\right.$$

It turns out that these sets, depicted in figure 2, are triangulations:

Proposition 6 *For any integer n greater than 2, and any relative integer k, $Z_n^-(k)$ and $Z_n^+(k)$ are triangulations of $\iota(n)$.*

Proof The result is proven by induction on n. First, if $n = 3$, the boundary edges on $\iota(n)$ are $\{0, 1\}$, $\{1, 2\}$, and $\{0, 2\}$. As there are no other edges on $\iota(n)$, this proves that $Z_n^-(k)$ and $Z_n^+(k)$ are both equal to $\beta(\iota(3))$ in this case. As $\beta(\iota(3))$ is a triangulation of $\iota(3)$, the result follows.

Now assume that $Z_n^-(k)$ and $Z_n^+(k)$ are triangulations of $\iota(n)$ for some integer n greater than 2, and any relative integer k. It is first shown that:

$$Z_{n+1}^+(-1) = Z_n^-(0) \cup \{\{0, n\}, \{n, n-1\}\}. \tag{7}$$

Let l and m be non-necessarily distinct elements of $\{-1, 0\}$. Consider an edge $\{i, j\}$ that belongs to $f_{n-m}(l+m)$. In this case, $i+j$ is congruent to $l+m$ modulo $n-m$. As the vertices of $\{i, j\}$ are distinct elements of $\iota(n-m)$, their sum satisfies:

$$1 \leq i + j \leq 2(n-m) - 3.$$

The only integer congruent to $l + m$ modulo $n - m$ in this range is $n + l$ which shows that $i + j = n + l$. Hence $i + j$ is congruent to $l - m - 1$ modulo $n+m+1$, showing that edge $\{i, j\}$ belongs to $f_{n+m+1}(l-m-1)$. One therefore obtains the following inclusion:

$$f_{n-m}(l + m) \subset f_{n+m+1}(l - m - 1).$$

Combining the four inclusions found when l and m take on any possible value in $\{-1, 0\}$ yields:

$$f_n(-1) \cup f_n(0) = f_{n+1}(-2) \cup f_{n+1}(-1). \tag{8}$$

Now observe that the only boundary edge of $\iota(n)$ that is not a boundary edge of $\iota(n+1)$, that is edge $\{0, n-1\}$, belongs to $f_n(-1)$. Hence, following the definition of sets $Z_n^-(0)$ and $Z_{n+1}^+(-1)$, equation (8) states that the first of these sets is a subset of the other. In addition, the only two boundary edges of $\iota(n+1)$ that are not a boundary edges of $\iota(n)$ are $\{0, n\}$ and $\{n-1, n\}$. Equation (7) therefore follows from (8). As $\{0, n\}$ and $\{n-1, n\}$ are boundary edges on $\iota(n+1)$, then they do not cross any edge on $\iota(n+1)$. Moreover, (7) states that $Z_{n+1}^+(-1)$ as exactly two more edges than $Z_n^-(0)$. By induction, $Z_n^-(0)$ is a triangulation and it follows from proposition 2 that $Z_{n+1}^+(0)$ is also a triangulation.

Finally, as for any relative integer k, $Z_{n+1}^-(k)$ and $Z_{n+1}^+(k)$ can be obtained from $Z_{n+1}^+(-1)$ by boundary-preserving permutations of $\iota(n+1)$, it follows from proposition 1 that these two sets are triangulations. □

Triangulations $Z_n^-(k)$ and $Z_n^+(k)$ are depicted in figure 2 depending on the parity of n. One can see that the interior edges of these triangulations are arranged as a zigzag that starts near the vertex of $\iota(n)$ congruent to k and ends near the vertex of $\iota(n)$ congruent to $k + \lfloor n/2 \rfloor$. Note, however, that the way this zigzag ends depends on the parity of n. In particular, if n is even then $Z_n^-(k)$ and $Z_n^+(k)$ respectively coincide with $Z_n^-(k + n/2)$ and $Z_n^+(k + n/2)$. If, on the other hand, n is odd, then $Z_n^-(k)$ and $Z_n^+(k)$ are respectively equal to $Z_n^+(k + \lfloor n/2 \rfloor)$ and to $Z_n^-(k + \lceil n/2 \rceil)$. This observation will be important for the construction of a maximally distant pair of triangulations.



Fig. 2 Triangulations $Z_n^-(k)$ (left) and $Z_n^+(k)$ (right), depicted depending on the parity of n. The interrupted edges show that the length of each zigzag is arbitrary.

6.2 A first pair of triangulations

Two triangulations are now built using two successive contractions will at the ends of zigzags. Each of these contractions will introduce one comb with either three or four teeth. Recall that contracting an edge in a triangulation decreases the number of its vertices. Hence, in order to build triangulations of $\iota(n)$, these contractions will be carried out in triangulations of $\iota(n+4)$. Let n be an integer greater than 2. Consider the two following edges on $\iota(n+4)$:

$$\varepsilon = \{\lfloor n/2 \rfloor - 1, \lfloor n/2 \rfloor + 1\} \text{ and } \varsigma = \{\lfloor n/2 \rfloor + 2, \lfloor n/2 \rfloor + 4\}.$$

One can define two triangulations W_n^- and W_n^+ of $\iota(n)$ by performing the following sequences of contractions in $Z_{n+4}^-(0)$ and in $Z_{n+4}^+(-1)$:

$$W_n^- = \varLambda_{\varepsilon}^{\star} \circ \varLambda_{\{n+1,n+3\}}^{\star}(Z_{n+4}^-(0)) \text{ and } W_n^+ = \varLambda_{\{0,2\}}^{\star} \circ \varLambda_{\varsigma}^{\star}(Z_{n+4}^-(-1)).$$

Observe that each of these contractions removes two vertices. Triangulations W_n^- and W_n^+ are depicted in figure 3 when n is greater than 8. It can be seen that, for such values of n, the contractions of ε and ς will produce two combs respectively at vertex $\lfloor n/2 \rfloor - 1$ in W_n^- and at vertex $\lfloor n/2 \rfloor$ in W_n^+ whose numbers of teeth (three or four) depend on the parity of n. The contractions of $\{0,2\}$ and $\{n+1, n+3\}$ always respectively result in a comb with four teeth at vertex 0 in W_n^+ and in a comb with three teeth at vertex $n-1$ in W_n^-. When n is greater than 9, the two combs contained in either of these triangulations are connected by a zigzag, and when $n = 9$, these two combs have a common tooth. When n is less than 9, the combs in W_n^- and W_n^+ loose teeth, or even disappear. Indeed, if $n = 3$, then W_n^- and W_n^+ are both equal to the unique triangulation of $\iota(3)$ made up of a single triangle. Triangulations W_n^- and W_n^+ are depicted in figure 4 when $4 \leq n \leq 8$. It will be shown in section 8 that W_n^- and W_n^+ have flip distance $2n - 10$ when n is greater than 12. Consider the pair made up of these two triangulations:

$$A_n = \{W_n^-, W_n^+\}.$$

While an accurate depiction of these triangulations requires a disjunction on the value of n, some properties of W_n^- and W_n^+ hold whenever n is greater



Fig. 3 Triangulations W_n^- (left) and W_n^+ (right), depicted when n is greater than 8. The interrupted edges show that the length of each zigzag is arbitrary.

than 4. A such property will now be used to obtain a first lower bound on $\delta(A_n)$. Assume that n is greater than 9 and call $\upsilon = \{n-2, n-1\}$. Now, consider triangulation W_n^-, shown in figure 3. Contracting edge υ in this triangulation maps vertex $n-1$ to vertex $n-2$. Hence, the triangulation $\Lambda_\upsilon^\star(W_n^-)$ that results from this contraction has a comb with four teeth at vertex $n-2$. Further observe that $\Lambda_\upsilon^\star(W_n^-)$ is the image of W_{n-1}^+ by the boundary-preserving permutation α of $\iota(n-1)$ that exchanges 0 and $n-2$:

$$\alpha : \begin{array}{l} \pi \to \pi, \\ i \mapsto n-2-i, \end{array}$$

where $\pi = \iota(n-1)$. Note in particular that α maps the comb at vertex $\lfloor (n-1)/2 \rfloor$ in triangulation W_{n-1}^+ to the comb at vertex $\lfloor n/2 \rfloor - 1$ in triangulation $\Lambda_\upsilon^\star(W_n^-)$.

Similarly, contracting edge υ in triangulation W_n^+ maps vertex $n-1$ to vertex $n-2$. The comb at vertex 0 therefore loses a tooth in the triangulation that results from this contraction. In other words, triangulation $\Lambda_\upsilon^\star(W_n^+)$ has a comb with three teeth at vertex 0. In fact, it can be seen in figure 3 that this triangulation is the image of W_{n-1}^- by α. Hence, when n is greater than 9,

$$\Lambda_\upsilon^\star(W_n^-) = \alpha(W_{n-1}^+) \text{ and } \Lambda_\upsilon^\star(W_n^+) = \alpha(W_{n-1}^-).$$

It can be checked using figure 4 that these equalities still hold when n ranges from 4 to 9. As a consequence,

Theorem 7 *Let n be an integer greater than 3. If $\vartheta_{\{n-2,n-1\}}(A_n)$ is not less than 2, then $\delta(A_n) \geq \delta(A_{n-1}) + 2$.*

Proof Denote $\upsilon = \{n-2, n-1\}$ and assume that $\vartheta_\upsilon(A_n) \geq 2$. According to corollary 1, the following inequality holds:

$$\delta(A_n) \geq \delta \circ \Lambda_\upsilon^\star(A_n) + 2. \tag{9}$$

It turns out that the right-hand side of this inequality is precisely equal to $\delta(A_{n-1})+2$. Indeed, consider the boundary-preserving permutation α of $\iota(n-1)$ that exchanges 0 and $n-2$. Recall that the two elements of pair A_{n-1}



Fig. 4 Triangulations W_n^- and W_n^+ when n ranges from 4 to 8.

are W_{n-1}^- and W_{n-1}^+. One can check using figures 3 and 4 that the images of these triangulations by α are respectively equal to the images of W_n^+ and W_n^- by $\Lambda_\upsilon^\star$. Hence, lemma 6 yields:

$$\delta \circ \Lambda_\upsilon^\star(A_n) = \delta(A_{n-1}). \tag{10}$$

The result is obtained by combining (9) and (10). □

Theorem 7 is a (small) step towards obtaining a lower bound of the form $2n + O(1)$ on the flip distance of pair A_n. Observe that the contraction used in its proof can be iterated in order to bound $\delta(A_n)$ below using $\delta(A_{n-k})$ for some positive integer k less than $n-2$. In particular, if n is greater than 4, then successively contracting edges $\{n-2, n-1\}$ and $\{0,1\}$ in triangulations W_n^- and W_n^+ directly results in respectively W_{n-2}^- and W_{n-2}^+ (see figures 3 and 4). Now observe that the order of these two contractions can be reversed without changing the result, provided edge $\{n-3, n-2\}$ is contracted instead of edge $\{n-2, n-1\}$ (recall that $\Lambda_{\{0,1\}}$ decreases by 1 all the positive integers in $\iota(n)$). In other words, contracting edge $\{0,1\}$ first and then edge $\{n-3, n-2\}$ in triangulations W_n^- and W_n^+ also respectively results in W_{n-2}^- and W_{n-2}^+. One obtains the following theorem as a consequence:

Theorem 8 *Let n be an integer greater than 7. If $\vartheta_{\{0,1\}}(A_n)$ is not less than 3, then $\delta(A_n) \geq \delta(A_{n-2}) + 4$.*

Proof Denote $\kappa = \{0,1\}$ and $\upsilon = \{n-3, n-2\}$. Assume that $\vartheta_\kappa(A_n) \geq 3$. According to corollary 1, the following inequality holds:

$$\delta(A_n) \geq \delta \circ \Lambda_\kappa^\star(A_n) + 3. \tag{11}$$

Recall that the two elements of pair A_n are W_n^- and W_n^+. One can see in figures 3 and 4 that the links of edge $\{n-2, n-1\}$ in W_n^- and W_n^+ are respectively $\{3\}$ and $\{0\}$ when n is greater than 7. When κ is contracted in triangulations W_n^- and W_n^+, every positive integer in $\iota(n)$ is mapped to its



Fig. 5 Triangulations Y_n^- (left) and Y_n^+ (right), depicted in solid lines depending on the parity of n. Triangulation X_n^-, obtained by flipping edge $\{4, 6\}$ in Y_n^- is sketched using dotted lines. The interrupted edges show that the length of each zigzag is arbitrary.

immediate predecessor. Hence, Λ_κ maps edge $\{n-2, n-1\}$ to υ. In addition, the link of υ in $\Lambda_\kappa^\star(W_n^+)$ is still $\{0\}$ and its link in $\Lambda_\kappa^\star(W_n^-)$ becomes $\{2\}$. As these two links are distinct, it follows from lemma 5 that $\vartheta_\upsilon \circ \Lambda_\kappa^\star(A_n)$ is positive and corollary 1 yields the following inequality:

$$\delta \circ \Lambda_\kappa^\star(A_n) \geq \delta \circ \Lambda_\upsilon^\star \circ \Lambda_\kappa^\star(A_n) + 1. \tag{12}$$

One can see in figures 3 and 4 that successively contracting κ and then υ in W_n^- and W_n^+ respectively results in triangulations W_{n-2}^- and W_{n-2}^+. As pair A_{n-2} is precisely made up of these two triangulations, the right-hand side of inequality (12) simplifies to $\delta(A_{n-2})+1$ and the result is immediately obtained by combining (11) and (12). □

6.3 Two more pairs of triangulations and a result on flip distances

If the conditions in the statements of theorem 7 (i.e. $\vartheta_{\{n-2,n-1\}}(A_n) \geq 2$) and theorem 8 (i.e. $\vartheta_{\{0,1\}}(A_n) \geq 3$) never fail together, then the flip distance of A_n admits a lower bound of the form $2n + O(1)$. No argument is available, though, to prove that these conditions are necessarily met. In order to work around this missing argument, it will be shown (theorem 12 in section 7) that a more elaborate recursive lower bound holds when n is greater than 12. This bound will be obtained from a careful study of minimal paths between the two triangulations of A_n under the assumption that $\vartheta_{\{n-2,n-1\}}(A_n) \leq 1$ and $\vartheta_{\{0,1\}}(A_n) \leq 2$. This study will rely on the two auxiliary pairs of triangulations B_n and C_n that are now introduced.

Assume that n is greater than 7. In this case, $\lceil n/2 \rceil + 7 \leq n + 3$. Hence, one can define two interior edges on $\iota(n+4)$ by:

$$\varepsilon = \{\lceil n/2 \rceil + 4, \lceil n/2 \rceil + 6\} \text{ and } \varsigma = \{\lceil n/2 \rceil + 5, \lceil n/2 \rceil + 7\}.$$

Two triangulations of $\iota(n)$ can be built by carrying out the following contractions in triangulations $Z_{n+4}^+(5)$ and $Z_{n+4}^+(2)$:

$$Y_n^- = \Lambda_{\{6,8\}}^\star \circ \Lambda_\varepsilon^\star(Z_{n+4}^+(5)) \text{ and } Y_n^+ = \Lambda_{\{3,5\}}^\star \circ \Lambda_\varsigma^\star(Z_{n+4}^+(2)).$$

Triangulations Y_n^- and Y_n^+ are shown in figure 5. In this figure, the dotted edges depict a variation of Y_n^- that will be introduced below. One can see that, when n is greater than 8, the two contractions carried out for the construction of Y_n^- introduce two combs at vertices 6 and $\lceil n/2 \rceil - 1$ connected by a zigzag. The first of these combs always has three teeth, and the other one has three or four teeth depending on the parity of n. If n is equal to 8, these two combs merge into a single one with five teeth at vertex 6, depicted in the left of figure 10. For any integer n greater than 7, one can see in the left of figure 5 that Y_n^+ admits two combs at vertices 3 and $\lceil n/2 \rceil + 3$. The comb at vertex 3 always has three teeth, and the other one has three or four teeth depending on the parity of n. It can further be seen in figures 3 and 5 that Y_n^+ is the image of W_n^- by a boundary-preserving map.

Observe that edge $\{4, 6\}$ always belongs to Y_n^-. The triangulation obtained by flipping this edge in Y_n^- will be denoted by X_n^-:

$$X_n^- = \mathfrak{F}(\{0, 2\}, Y_n^-).$$

Hence, triangulation X_n^- differs from Y_n^- by exactly one edge depicted as a dotted line in figures 5 and 10. For any integer n greater than 8, consider the two following pairs of triangulations of $\iota(n)$:

$$B_n = \{Y_n^-, Y_n^+\} \text{ and } C_n = \{X_n^-, Y_n^+\}.$$

It is now shown that, if the conditions on $\vartheta_{\{n-2,n-1\}}(A_n)$ and $\vartheta_{\{0,1\}}(A_n)$ in the statements of theorems 7 and 8 fail together, and if n is greater than 8, then there always exists a pair P of triangulations of $\iota(n-1)$ so that $\delta(A_n)$ is not less that $\delta(P) + 3$. According to the following theorem, it can be required that pair P is either equal to B_{n-1} or to C_{n-1} and that the image of P by $\vartheta_{\{3,4\}}$ is not greater than 1:

Theorem 9 *Let n be an integer greater than* 8*. If the images of A_n by maps $\vartheta_{\{n-2,n-1\}}$ and $\vartheta_{\{0,1\}}$ are less than respectively* 2 *and* 3*, then there exists $P \in \{B_{n-1}, C_{n-1}\}$ so that $\delta(A_n) = \delta(P) + 3$ and $\vartheta_{\{3,4\}}(P)$ is less than* 2*.*

Proof Denote $\kappa = \{0, 1\}$ and $\upsilon = \{n - 2, n - 1\}$. Assume that $\vartheta_\kappa(A_n) \leq 2$ and that $\vartheta_\upsilon(A_n) \leq 1$. Consider a minimal path ψ of length k from W_n^- to W_n^+. Since $\vartheta_\upsilon(A_n) \leq 1$, set $\gamma_\upsilon(\psi)$ contains at most one element. In fact, it contains exactly one element. Indeed, as n is greater than 8, the links of υ in triangulations W_n^- and W_n^+ are respectively $\{3\}$ and $\{0\}$ (see figure 3). Since these two links are distinct, it follows from lemma 5 that $\gamma_\upsilon(\psi)$ is non-empty. Hence, it admits exactly one element j. In this case, according to theorem 5, edge $\{0, 3\}$ belongs to $\psi(j)$ and to $\psi(j + 1)$. Moreover, as all the boundary edges of quadrilateral $\{0, 1, 2, 3\}$ other than $\{0, 3\}$ are also boundary edge of $\psi(j)$, then the following inclusion holds:

$$\beta(\{0, 1, 2, 3\}) \subset \psi(j).$$

As a consequence, $\psi(j)$ necessarily contains exactly one diagonal of quadrilateral $\{0, 1, 2, 3\}$ (i.e. edge $\{0, 2\}$ or edge $\{1, 3\}$). Respectively denote by ε and



Fig. 6 Left: the triangulation obtained by successively flipping edges $\{2, n-1\}$, $\{1, n-1\}$, and $\{3, n-1\}$ in W_n^-, corresponding to the case $\varepsilon = \{1, 3\}$ in the proof of theorem 9. Right: the triangulation obtained by successively flipping edges $\{1, n-1\}$, $\{2, n-1\}$, and $\{3, n-1\}$ in W_n^-, corresponding to the case $\varepsilon = \{0, 2\}$ in the same proof.

ε' these two diagonals with the convention that $\varepsilon \in \psi(j)$ and $\varepsilon' \notin \psi(j)$. It can be seen in figure 3 that ε and ε' are the links of edges $\{1, n-1\}$ and $\{2, n-1\}$ in triangulation W_n^-. Denote by ς and ς' these two edges of W_n^- with the requirement that ε and ε' are the respective links of ς and ς' in W_n^-.

Consider edges ε, $\{0, 3\}$, and $\{0, n-2\}$. One can see in figure 3 that these three edges are precisely introduced in this order when successively flipping ς, ς', and $\{3, n-1\}$ in W_n^-. It has been shown in the preceding paragraphs that ε and $\{0, 3\}$ respectively belong to triangulations $\psi(j)$ and $\psi(j+1)$. In addition, $\{0, n-2\}$ is an edge of $\psi(k)$. As $j \leq j+1 \leq k$, then according to theorem 3, it can be assumed without loss of generality that ς, ς', and $\{3, n-1\}$ are the first three edges flipped in this order along path ψ. In other words,

$$\psi(1) = \mathfrak{F}(\varsigma, \psi(0)),\ \psi(2) = \mathfrak{F}(\varsigma', \psi(1)), \text{ and } \psi(3) = \mathfrak{F}(\{3, n-1\}, \psi(2)).$$

Triangulation $\psi(3)$ is shown in figure 6 for the two possible values of ε. According to the construction of $\psi(1)$, $\psi(2)$, and $\psi(3)$, j is equal to 2. In addition, $\gamma_\kappa(\psi)$ either contains 0 (if $\varepsilon = \{0, 2\}$) or 1 (if $\varepsilon = \{1, 3\}$). As a consequence $\gamma_\kappa(\psi)$ contains an element of $\iota(3)$. As in addition, $\vartheta_\kappa(A_n)$ is not greater than 2, then $\gamma_\kappa(\psi)$ contains at most one element of $\iota(k) \setminus \iota(3)$. Further observe that one can prescribe the triangulations visited by path ψ between $\psi(3)$ and W_n^+ to be those of an arbitrarily chosen minimal path from $\psi(3)$ to W_n^+. As $\gamma_\kappa(\psi)$ contains at most one element greater than 2, the following inequality is obtained as a consequence:

$$\vartheta_\kappa(\{\psi(3), W_n^+\}) \leq 1 \tag{13}$$

Now consider the boundary-preserving permutation α of $\iota(n-1)$ that decreases by 3 the elements of $\iota(n-1) \setminus \iota(3)$. In other words, α is defined as follows on every integer i in $\iota(n-1)$:

$$\alpha(i) = \begin{cases} n - 4 + i & \text{if } 0 \leq i \leq 2, \\ i - 3 & \text{otherwise.} \end{cases}$$

It can be seen in figures 3 and 5 that $\alpha(Y_{n-1}^{+})$ is a subset of W_n^{+} and that the elements of $W_n^{+} \setminus \alpha(Y_{n-1}^{+})$ are edges $\{0, n-1\}$ and $\{n-2, n-1\}$. Call:

$$U = \begin{cases} Y_{n-1}^{-} & \text{if } \varepsilon = \{1,3\}, \\ X_{n-1}^{-} & \text{if } \varepsilon = \{0,2\}. \end{cases}$$

One can see in figures 5 and 6 that in both cases, the image of U by α is a subset of $\psi(3)$, and that the set obtained by removing the elements of $\alpha(U)$ from $\psi(3)$ is made up of edges $\{0, n-1\}$ and $\{n-2, n-1\}$ (dotted in figure 6). Further observe that α maps $\{3,4\}$ to κ. Hence, denoting by P the pair $\{U, Y_{n-1}^{+}\}$, it follows from lemma 6, that:

$$\delta(\{\psi(3), W_n^{+}\}) = \delta(P) \text{ and } \vartheta_{\kappa}(\{\psi(3), W_n^{+}\}) = \vartheta_{\{3,4\}}(P).$$

As ψ is a minimal path from W_n^{-} to W_n^{+}, the left-hand side of the first equality is $\delta(A_n) - 3$, proving that $\delta(A_n) = \delta(P) + 3$. Combining the second equality with (13), one obtains $\vartheta_{\{3,4\}}(P) \leq 1$. Since P is either equal to B_{n-1} (if $\varepsilon = \{1,3\}$) or to C_{n-1} (if $\varepsilon = \{0,2\}$), the theorem is proven. □

7 A recursive lower bound on $\delta(A_n)$

The conditions on $\vartheta_{\{n-2,n-1\}}(A_n)$ and on $\vartheta_{\{0,1\}}(A_n)$ in theorems 7, 8, and 9 are complementary. Combining the inequalities provided by these theorems, one therefore obtains the following lower bound on $\delta(A_n)$:

Proposition 7 *For any integer n greater than 8,*

$$\delta(A_n) \geq \min(\{\delta(A_{n-1}) + 2, \delta(A_{n-2}) + 4, \delta(B_{n-1}) + 3, \delta(C_{n-1}) + 3\}).$$

This inequality is improved in this section into a recursive lower bound on $\delta(A_n)$, obtained bounding $\delta(B_n)$ and $\delta(C_n)$ below by respectively $\delta(A_{n-5}) + 9$ and $\delta(A_{n-4}) + 7$ whenever n is greater than 11. These two bounds will be obtained under the respective assumptions that the images of B_n and C_n by $\vartheta_{\{3,4\}}$ are less than 2. Note, though, that according to the conclusion of theorem 9, these assumptions are not restrictive.

7.1 A lower bound on the flip distance of B_n

The bound on $\delta(B_n)$ is first obtained by invoking theorem 6 for the successive contractions of edges $\{4,6\}$ and $\{0,3\}$. The following lemma corresponds to the first of these contractions:

Lemma 7 *Let n be an integer greater than 11. If $\vartheta_{\{3,4\}}(B_n)$ is less than 2, then $\delta(B_n) \geq \delta \circ \Lambda_{\{4,6\}}^{\star}(B_n) + 4$.*

Proof Assume that $\vartheta_{\{3,4\}}(B_n) \leq 1$ and recall that the two elements of B_n are Y_n^- and Y_n^+. Let ψ be a minimal path from Y_n^- to Y_n^+. As $\vartheta_{\{3,4\}}(B_n) \leq 1$, then $\gamma_{\{3,4\}}(\psi)$ contains at most one element. As can be seen in figure 5, the links of $\{3,4\}$ in triangulations Y_n^- and Y_n^+ are respectively $\{6\}$ and $\{n-1\}$. As n is greater than 11, these links are distinct and it follows from lemma 5 that $\gamma_{\{3,4\}}(\psi)$ contains exactly one element, say j. Denote:

$$P = \{Y_n^-, \psi(j+1)\} \text{ and } Q = \{\psi(j+1), Y_n^+\}.$$

According to the triangular inequality:

$$\delta \circ \Lambda^\star_{\{4,6\}}(P) + \delta \circ \Lambda^\star_{\{4,6\}}(Q) \geq \delta \circ \Lambda^\star_{\{4,6\}}(B_n). \tag{14}$$

Observe that the restriction of ψ to $\iota(j+2)$ is a minimal path between the two triangulations in pair P. As j is the only element of $\gamma_{\{3,4\}}(\psi)$, it then follows from the two first assertions in the statement of theorem 5 that edges $\{3,6\}$ and $\{4,n-1\}$ respectively belong to $\psi(j)$ and $\psi(j+1)$. As these edges are crossing, and as $\psi(j)$ and $\psi(j+1)$ are adjacent, then, $\psi(j+1)$ is obtained replacing edge $\{3,6\}$ in $\psi(j)$ by edge $\{4,n-1\}$. Further observe that quadrilateral $\{3,4,6,n-1\}$ contains the two vertices of edge $\{4,6\}$. This shows that j belongs to $\gamma_{\{4,6\}}(\psi)$ and therefore that $\vartheta_{\{4,6\}}(P) \geq 1$. One then obtains the following inequality from corollary 1:

$$\delta(P) \geq \delta \circ \Lambda^\star_{\{4,6\}}(P) + 1. \tag{15}$$

Recall that $\psi(j+1)$ is obtained replacing edge $\{3,6\}$ in $\psi(j)$ by edge $\{4,n-1\}$. Hence, all the boundary edges of quadrilateral $\{3,4,6,n-1\}$ as well as its diagonal $\{4,n-1\}$ belong to $\psi(j+1)$. In particular, the link of $\{4,6\}$ in $\psi(j+1)$ is $\{5,n-1\}$. Now consider the comb at vertex $\lceil n/2 \rceil + 3$ in triangulation Y_n^+, depicted in the right of figure 5. As n is greater than 11, then the following inequality holds:

$$\lceil n/2 \rceil + 3 \leq n - 3.$$

One can see in figure 5 that, in this case, the links of edges $\{4,5\}$ and $\{5,6\}$ in triangulation Y_n^+ are respectively $\{n-2\}$ and $\{n-3\}$. This shows that $\{4,5\}$ and $\{5,6\}$ have distinct links in Y_n^+ and that these two links are also disjoint from that of edge $\{4,6\}$ in triangulation $\psi(j+1)$. Hence, the conditions of theorem 6 are satisfied if one respectively replaces U, V, and κ by Y_n^+, $\psi(j+1)$, and $\{4,6\}$ in the statement of this theorem. In this case, $\overline{\kappa}$ contains 3 vertices of $\iota(n)$ and the following inequality is found as a consequence:

$$\delta(Q) \geq \delta \circ \Lambda^\star_{\{4,6\}}(Q) + 3. \tag{16}$$

Since ψ is a minimal path, it follows from the definition of pairs P and Q that $\delta(P) + \delta(Q)$ is equal to $\delta(B_n)$. The result is therefore obtained by combining inequalities (14), (15), and (16). □



Fig. 7 The images of triangulations Y_n^- when n is greater than 11 (left) and Y_n^+ when n is greater than 15 (right) by map $\varLambda^\star_{\{4,6\}}$, depicted depending on the parity of n.

Denote $\upsilon = \{4,6\}$. The images of Y_n^- and Y_n^+ by $\varLambda^\star_\upsilon$ are depicted in figures 7 and 8. Note that $\varLambda^\star_\upsilon(Y_n^-)$ and $\varLambda^\star_\upsilon(Y_n^+)$ are triangulations of $\iota(n-2)$, because contracting υ in $\iota(n)$ removes two vertices from this polygon. In particular, the greatest vertex of these two triangulations is $n-3$. One can see that $\varLambda^\star_\upsilon(Y_n^-)$ is made up of a zigzag that starts at vertex 4 and ends at vertex $\lceil n/2\rceil$ with a comb whose number of teeth depends on the parity of n.

While the sketch of $\varLambda^\star_\upsilon(Y_n^-)$ in figure 7 precisely shows how this triangulation is made for any integer n greater than 11, that of $\varLambda^\star_\upsilon(Y_n^+)$ in the same figure can only be used when n is greater than 15. Indeed, this sketch shows that, for these values of n, triangulation $\varLambda^\star_\upsilon(Y_n^+)$ contains three combs at vertices 3, 4, and $\lceil n/2\rceil+1$ and that the two latter combs are connected by a zigzag. If $12 \leq n \leq 15$, figure 8 shows that this zigzag disappears as edge $\{4, \lceil n/2\rceil+1\}$ becomes a common tooth of the two combs at its endpoints. The comb at vertex $\lceil n/2\rceil+1$ even reduces to a simple zigzag when $n=12$.

Invoking theorem 6 for the contraction of edge $\{0,3\}$ in the triangulations of pair $\varLambda^\star_\upsilon(B_n)$, one obtains the following inequality:

Lemma 8 *For any integer n greater than* 11*,*

$$\delta\circ\varLambda^\star_{\{4,6\}}(B_n) \geq \delta\circ\varLambda^\star_{\{0,3\}}\circ\varLambda^\star_{\{4,6\}}(B_n)+5.$$

Proof Consider an integer n greater than 11, and denote $\upsilon=\{4,6\}$. One can see in figure 7 when $n\geq 16$ and in figure 8 when $12\leq n\leq 15$ that the link of $\{0,3\}$ in $\varLambda^\star_\upsilon(Y_n^+)$ is always $\{1,n-3\}$ (recall that $\varLambda^\star_\upsilon(Y_n^+)$ is a triangulation of $\iota(n-2)$, and that its greatest vertex is therefore $n-3$).

Now consider the comb at vertex $\lceil n/2\rceil$ in triangulation $\varLambda^\star_\upsilon(Y_n^-)$, shown in the left of figure 7. Since n is greater than 11, $\lceil n/2\rceil$ is not less than 6. Hence, one can see in this figure that, for such values of n, the links of edges $\{0,1\}$, $\{1,2\}$, and $\{2,3\}$ in triangulation $\varLambda^\star_\upsilon(Y_n^-)$ are respectively $\{6\}$, $\{5\}$, and $\{4\}$. In particular, these three links are pairwise disjoint.

As n is greater than 11, the links of edges $\{0,1\}$, $\{1,2\}$, and $\{2,3\}$ in triangulation $\varLambda^\star_\upsilon(Y_n^-)$ are also disjoint from that of $\{0,3\}$ in $\varLambda^\star_\upsilon(Y_n^+)$. Hence, the conditions of theorem 6 are satisfied if one respectively replaces U, V, and


$\Lambda^\star_{\{4,6\}}(Y^+_{12})$ $\Lambda^\star_{\{4,6\}}(Y^+_{13})$ $\Lambda^\star_{\{4,6\}}(Y^+_{14})$ $\Lambda^\star_{\{4,6\}}(Y^+_{15})$

Fig. 8 The images of triangulations Y^+_n by map $\Lambda^\star_{\{4,6\}}$ when $12 \leq n \leq 15$.

κ by $\Lambda^\star_\upsilon(Y^-_n)$, $\Lambda^\star_\upsilon(Y^+_n)$, and $\{0,3\}$ in the statement of this theorem. In this case, $\overline{\kappa}$ contains four vertices of $\iota(n)$ and theorem 6 yields:

$$\delta \circ \Lambda^\star_\upsilon(B_n) \geq \delta \circ \Lambda^\star_{\{0,3\}} \circ \Lambda^\star_\upsilon(B_n) + 5.$$

Since $\upsilon = \{4,6\}$, this completes the proof. □

Let n be an integer greater than 11. Denote $\kappa = \{0,3\}$ and $\upsilon = \{4,6\}$. One can see in figures 7 (when $n \geq 16$) and 8 (when $12 \leq n \leq 15$) that contracting edge κ in $\Lambda^\star_\upsilon(Y^+_n)$ removes one of the combs from this triangulation, and leaves the other combs unaffected, except for the numbering of their vertices. In particular, $\Lambda^\star_\kappa \circ \Lambda^\star_\upsilon(Y^+_n)$ admits a comb at vertex 1. This comb has four teeth when $n \geq 14$ and three teeth when $12 \leq n \leq 13$. Note that $\{0, 1, n-6\}$ is an ear of this triangulation. Further observe that, when n is greater than 12, triangulation $\Lambda^\star_\kappa \circ \Lambda^\star_\upsilon(Y^+_n)$ has another comb at vertex $\lceil n/2 \rceil - 2$. This comb has three teeth when $n = 13$, and three or four teeth depending on the cardinality of n when n is greater than 13. If $n = 12$, the comb at vertex $\lceil n/2 \rceil - 2$ simplifies to a simple zigzag. In fact, it can be seen in these figures that triangulation $\Lambda^\star_\kappa \circ \Lambda^\star_\upsilon(Y^+_n)$ is the image of W^+_{n-5} by the boundary-preserving permutation α of $\iota(n-2)$ that subtracts 1 from the positive elements of $\iota(n-5)$.

Now consider triangulation $\Lambda^\star_\upsilon(Y^-_n)$ shown in the left of figure 7. Contracting edge κ in this triangulation introduces a new comb with three teeth at vertex 0. The comb already found in $\Lambda^\star_\upsilon(Y^-_n)$ is unaffected by this contraction except for the numbering of its vertices. In fact, one can see in these figures that $\Lambda^\star_\kappa \circ \Lambda^\star_\upsilon(Y^-_n)$ is the image of W^-_{n-5} by α. The following theorem is obtained as a consequence of these observations:

Theorem 10 *Let n be an integer greater than* 11. *If $\vartheta_{\{3,4\}}(B_n)$ is less than* 2 *then $\delta(B_n) \geq \delta(A_{n-5}) + 9$.*

Proof Assume that $\vartheta_{\{3,4\}}(B_n) \leq 1$. Respectively denote edges $\{0,3\}$ and $\{4,6\}$ by κ and υ. As n is greater than 11, one obtains the following inequality combining the results of lemmas 7 and 8:

$$\delta(B_n) \geq \delta \circ \Lambda^\star_\kappa \circ \Lambda^\star_\upsilon(B_n) + 9. \tag{17}$$

Consider the boundary-preserving permutation α of $\iota(n-5)$ that subtracts 1 from the positive elements of $\iota(n-5)$. In other words, α maps 0 to $n-6$ and every other element of $\iota(n-5)$ to its immediate predecessor. It can be seen in figures 7 and 8 that, since n is greater than 11, the images of $\Lambda_{\upsilon}^{\star}(Y_n^-)$ and $\Lambda_{\upsilon}^{\star}(Y_n^+)$ by $\Lambda_{\kappa}^{\star}$ respectively coincide with $\alpha(W_{n-5}^-)$ and $\alpha(W_{n-5}^+)$. Hence, according to lemma 6 and to the definition of pair A_{n-5},

$$\delta \circ \Lambda_{\kappa}^{\star} \circ \Lambda_{\upsilon}^{\star}(B_n) = \delta(A_{n-5}).$$

Combining this equality with inequality (17) completes the proof. □

7.2 A lower bound on the flip distance of C_n

The lower bound $\delta(C_n) \geq \delta(A_{n-4}) + 7$ is now established using the successive contractions of edges $\{4, 5\}$ and $\{0, 3\}$. The following lemma deals with the first of these contractions. Its proof is similar to that of lemma 7. In this case, though, corollary 3 can be invoked instead of theorem 6, which results in a substantial simplification.

Lemma 9 *Let n be an integer greater than 9. If $\vartheta_{\{3,4\}}(C_n)$ is less than 2, then $\delta(C_n) \geq \delta \circ \Lambda_{\{4,5\}}^{\star}(C_n) + 2$.*

Proof Assume that $\vartheta_{\{3,4\}}(C_n) \leq 1$ and recall that the two elements of C_n are X_n^- and Y_n^+. It can be seen in figure 5 that, for any integer n is greater than 9, edge $\{3, 5\}$ (dotted in the figure) belongs to triangulation X_n^-. Note that triangulations X_{10}^- and X_{11}^- are also depicted in a more precise way in figure 10. Further observe that Y_n^+ admits a comb at vertex $\lceil n/2 \rceil + 3$, and that the following inequality holds because n is greater than 9:

$$\lceil n/2 \rceil + 3 \leq n - 2.$$

Hence, as can be seen in figure 5 (and also in figure 10 when $10 \leq n \leq 11$), the links of edges $\{3, 4\}$ and $\{4, 5\}$ in triangulation Y_n^+ are respectively $\{n-1\}$ and $\{n-2\}$. In particular, these links are distinct. In this case, corollary 3 states that $\vartheta_{\{3,4\}}(C_n)$ and $\vartheta_{\{4,5\}}(C_n)$ cannot both be less than 2. By hypothesis, $\vartheta_{\{3,4\}}(C_n)$ is less than 2, showing that $\vartheta_{\{4,5\}}(C_n) \geq 2$. The desired inequality then immediately follows from corollary 1. □

Denote edge $\{4, 5\}$ by υ. The image of triangulation X_n^- by $\Lambda_{\upsilon}^{\star}$ is depicted in the left of figure 9 for any integer n greater than 7. One can see in this figure that, when n is greater than 8, $\Lambda_{\upsilon}^{\star}(X_n^-)$ is made up of a zigzag that starts at vertex 3 and ends at vertex $\lceil n/2 \rceil + 1$ with a comb whose number of teeth (three or four) depends on the parity of n. If $n = 8$, the zigzag disappears and $\Lambda_{\upsilon}^{\star}(X_n^-)$ is reduced to a comb with four vertices at vertex 5.

The image of triangulation Y_n^+ by $\Lambda_{\upsilon}^{\star}$ is shown in the right of figure 9 when n is greater than 11. It can be seen in this figure that, for such values of n, $\Lambda_{\upsilon}^{\star}(Y_n^+)$ admits three combs at vertices 3, 4, and $\lceil n/2 \rceil + 2$. The first two combs have 3 teeth each, and the third one has three or four teeth depending on the parity of n. In addition the combs at vertices 4 and $\lceil n/2 \rceil + 2$ are connected



Fig. 9 The images of triangulations X_n^- when n is greater than 7 (left) and Y_n^+ when n is greater than 11 (right) by map $\Lambda^\star_{\{4,5\}}$, depicted depending on the parity of n.

by a zigzag when n is greater than 13, and admit edge $\{4, \lceil n/2\rceil + 2\}$ as a common teeth when $12 \leq n \leq 13$.

Invoking theorem 6 for the contraction of edge $\{0, 3\}$ in pair $\Lambda^\star_\upsilon(C_n)$ provides the following result:

Lemma 10 *For any integer n greater than* 11*,*

$$\delta \circ \Lambda^\star_{\{4,5\}}(C_n) \geq \delta \circ \Lambda^\star_{\{0,3\}} \circ \Lambda^\star_{\{4,5\}}(C_n) + 5.$$

Proof Consider an integer n greater than 11 and denote edge $\{4, 5\}$ by υ. One can see in figure 9 that $\{0, 3\}$ is an edge of $\Lambda^\star_\upsilon(Y_n^+)$ whose link in this triangulation is $\{1, n-2\}$ (recall that $\Lambda^\star_\upsilon(Y_n^+)$ is a triangulation of $\iota(n-1)$, and that its greatest vertex is $n - 2$). Further observe that triangulation $\Lambda^\star_\upsilon(X_n^-)$ has a comb at vertex $\lceil n/2\rceil + 1$. As n is greater than 11, then:

$$\lceil n/2\rceil + 1 \geq 7.$$

As can be seen in figure 9, it follows from this inequality that the links of edges $\{0, 1\}$, $\{1, 2\}$, and $\{2, 3\}$ in triangulation $\Lambda^\star_\upsilon(X_n^-)$ are respectively $\{7\}$, $\{6\}$, and $\{5\}$. In particular, these three links are pairwise disjoint and, since n is greater than 11, they are also disjoint from that of $\{0, 3\}$ in $\Lambda^\star_\upsilon(Y_n^+)$. Hence, the conditions of theorem 6 are satisfied if one respectively replaces U, V, and κ by $\Lambda^\star_\upsilon(X_n^-)$, $\Lambda^\star_\upsilon(Y_n^+)$, and $\{0, 3\}$ in the statement of this theorem. In this case, $\overline{\kappa}$ contains four vertices of $\iota(n-1)$ and theorem 6 yields:

$$\delta \circ \Lambda^\star_\upsilon(C_n) \geq \delta \circ \Lambda^\star_{\{0,3\}} \circ \Lambda^\star_\upsilon(C_n) + 5.$$

Since $\upsilon = \{4, 5\}$, this completes the proof. □

Let n be an integer greater than 11. Respectively denote edges $\{0, 3\}$ and $\{4, 5\}$ by υ and κ. It can be seen in figure 9 that contracting κ in $\Lambda^\star_\upsilon(Y_n^+)$ removes one comb from this triangulation and leaves the other two combs unaffected, except for the numbering of their vertices. In particular, $\Lambda^\star_\kappa \circ \Lambda^\star_\upsilon(Y_n^+)$ has a comb with three teeth at vertex 0 and a comb at vertex $\lceil n/2\rceil - 1$ whose number of teeth is three or four depending on the parity of n. One can see using figure 3 that this $\varLambda_{\kappa}^{\star} \circ \varLambda_{\upsilon}^{\star}(Y_n^+)$ is in fact the image of W_{n-4}^- by the boundary-preserving permutation α of $\iota(n-4)$ that exchanges 0 and $n-5$:

$$\alpha : \begin{array}{l} \pi \rightarrow \pi, \\ i \mapsto n-5-i, \end{array}$$

where $\pi = \iota(n-4)$. Now consider triangulation $\varLambda_{\upsilon}^{\star}(X_n^-)$ depicted in the left of figure 9. It can be seen that contracting κ in this triangulation introduces a new comb with four teeth at vertex 0. Moreover, the other comb in $\varLambda_{\upsilon}^{\star}(X_n^-)$ is unaffected by this contraction, except for the numbering of its vertices. In particular, $\varLambda_{\kappa}^{\star} \circ \varLambda_{\upsilon}^{\star}(X_n^-)$ has a second comb at vertex $\lceil n/2 \rceil - 2$ with three or four teeth, depending on the parity of n. It turns out that $\varLambda_{\kappa}^{\star} \circ \varLambda_{\upsilon}^{\star}(X_n^-)$ is the image of W_{n-4}^+ by permutation α. Because of these observations, one obtains the following theorem from lemmas 9 and 10:

Theorem 11 *Let n be an integer greater than* 11. *If $\vartheta_{\{3,4\}}(C_n)$ is less than* 2 *then $\delta(C_n) \geq \delta(A_{n-4}) + 7$.*

Proof Assume that $\vartheta_{\{3,4\}}(C_n) \leq 1$. Respectively denote edges $\{0,3\}$ and $\{4,5\}$ by κ and υ. As n is greater than 11, one obtains the following inequality by invoking lemmas 9 and 10:

$$\delta(C_n) \geq \delta \circ \varLambda_{\kappa}^{\star} \circ \varLambda_{\upsilon}^{\star}(C_n) + 7. \tag{18}$$

Consider the boundary-preserving permutation of $\iota(n-4)$ that exchanges 0 and $n-5$. It can be seen, using figures 3 and 9 that the images of $\varLambda_{\upsilon}^{\star}(X_n^-)$ and $\varLambda_{\upsilon}^{\star}(Y_n^+)$ by $\varLambda_{\kappa}^{\star}$ respectively coincide with $\alpha(W_{n-4}^+)$ and $\alpha(W_{n-4}^-)$. Hence, according to lemma 6 and to the definition of pair A_{n-4},

$$\delta \circ \varLambda_{\kappa}^{\star} \circ \varLambda_{\upsilon}^{\star}(C_n) = \delta(A_{n-4}).$$

Combining this equality with (18) yields the desired inequality. □

7.3 A recursive lower bound on the flip distance of A_n

All the results needed to obtain an adequate recursive lower bound on $\delta(A_n)$ have been stated. This bound is found invoking theorems 7 to 11:

Theorem 12 *For any integer n greater than* 12*,*

$$\delta(A_n) \geq \min(\delta(A_{n-1}) + 2, \delta(A_{n-2}) + 4, \delta(A_{n-5}) + 10, \delta(A_{n-6}) + 12).$$

Proof Let n be an integer greater than 12. If $\vartheta_{\{n-2,n-1\}}(A_n)$ is not less than 2 or if $\vartheta_{\{0,1\}}(A_n)$ is not less than 3, the desired inequality follows either from theorem 7 or from theorem 8. Assume that $\vartheta_{\{n-2,n-1\}}(A_n) \leq 1$ and that $\vartheta_{\{0,1\}}(A_n) \leq 2$. According to theorem 9, there exists a pair of triangulations $P \in \{B_{n-1}, C_{n-1}\}$ so that the following inequality holds:

$$\delta(A_n) \geq \delta(P) + 3. \tag{19}$$

According to the same theorem, one can also require that $\vartheta_{\{3,4\}}(P)$ is not greater than 1. Under this condition, and because $n-1$ is greater than 11, theorems 10 and 11 state that $\delta(P)$ is not less than both $\delta(A_{n-6})+9$ and $\delta(A_{n-5})+7$. Hence, the desired inequality follows from (19). □

8 The diameters of associahedra

One of the consequences of theorem 12 is the existence of an integer k so that $\delta(A_n) \geq 2n-k$ is bounded below by a constant integer when n is greater than 12. It is shown in this section that 10 is a feasible value of integer k. According to lemma 1 and to theorem 2, this will settle the two conjectures on the diameter of associahedra.

As already mentioned, W_3^- and W_3^+ are both equal to the unique triangulation of $\iota(3)$, made up of a single triangle. One therefore obtains $\delta(A_3)=0$. Observe that this flip distance is already larger than $2n-10$ when one replaces n by 3. Consider pairs A_n shown in figure 4 when $4 \leq n \leq 8$. It can be seen that, if n is equal to 4 or to 5, then one can change W_n^- into W_n^+ using a sequence of flips so that each of the flips in this sequence introduces an edge of W_n^+. Hence, it follows from lemma 2 that $\delta(A_n)$ is equal to the number of interior edges of W_n^+ for these values of n, that is $\delta(A_n)=n-3$. Here, again, the flip distance of A_n is not less than $2n-10$.

Using more elaborate arguments further provides $\delta(A_n)$ for $6 \leq n \leq 8$:

Proposition 8 *The flip distances of pairs* A_3*,* A_4*,* A_5*,* A_6*,* A_7*, and* A_8 *are respectively* 0*,* 1*,* 2*,* 4*,* 5*, and* 7*.*

Proof The flip distances of A_3, A_4, and A_5 have already been obtained above. Consider the pair A_6, shown in figure 4. It is well known that this pair has flip distance 4. Indeed, the flip-graph of $\iota(6)$ has only 14 vertices and can easily be drawn (see for instance figure 1). The flip distance of A_6 can be obtained alternatively using a few simple arguments. Indeed, one can see that none of the flips that can be carried out in triangulation W_6^- will introduce an edge of W_6^+. Hence the second triangulation in any path from W_6^- to W_6^+ never shares an interior edge with W_6^+. From this triangulation, at least 3 other flips are needed because all three of its interior edges still have to be removed. This shows that $\delta(A_6) \geq 4$. The opposite inequality is obtained by exhibiting a path of length 4 between W_6^- and W_6^+. For instance, successively flipping edges $\{2,4\}$, $\{2,5\}$, $\{1,5\}$, and $\{3,5\}$ in W_6^- will result in W_6^+.

The flip distances of A_7 and A_8 can be easily deduced from that of A_6. Indeed, as edge $\{5,6\}$ has distinct links in W_7^- and W_7^+, it follows from lemma 5 that $\vartheta_{\{5,6\}}(A_7) \geq 1$, and then from corollary 1 that:

$$\delta(A_7) \geq \delta \circ \varLambda^{\star}_{\{5,6\}}(A_7) + 1.$$

It can be seen in figure 4 that contracting edge $\{5,6\}$ in W_7^- and in W_7^+ respectively results in triangulations W_6^- and W_6^+. In particular, the above inequality simplifies to $\delta(A_7) \geq \delta(A_6)+1$. It immediately follows that the flip distance of A_6 is not less than 5. As successively flipping edges $\{1,6\}$, $\{2,6\}$, $\{2,5\}$, $\{2,4\}$, and $\{0,2\}$ transforms W_7^- into W_7^+, the opposite inequality also holds, proving the the flip distance of A_7 is indeed 5.

Now consider pair A_8 shown in the right of figure 4. One can see that edge $\{3,7\}$ crosses both $\{0,6\}$ and $\{2,4\}$. Hence, the conditions of corollary 2 hold when U, V, κ, υ, x, and y are are respectively replaced by W_8^+, W_8^-, $\{2,3\}$, $\{6,7\}$, 2, and 6. In particular, in this case, $\lambda(\upsilon,U)$ and $\{x\}\cup\lambda(\kappa,U)$ are equal to $\{0\}$ and $\{2,4\}$. Hence the first of these two sets is not a subset of the other one, and the third assertion in the statement of corollary 2 holds. One therefore obtains from this corollary that $\vartheta_{\{2,3\}}(A_8)$ and $\vartheta_{\{6,7\}}(A_8)$ cannot both be less than 2. Hence, corollary 1 yields:

$$\delta(A_8) \geq \delta \circ \varLambda_\kappa^\star(A_8)+2,$$

where κ is equal to either $\{2,3\}$ or $\{6,7\}$. In both cases, one can find a boundary-preserving permutation of $\iota(7)$ that maps $\varLambda_\kappa^\star(A_8)$ to A_7. Therefore, following lemma 6, $\delta(A_8) \geq \delta(A_7)+2$, which shows that the flip distance of A_7 is not less than 7. Finally, sequentially flipping the following seven edges transforms W_8^- into W_8^+:

$$\{1,7\},\ \{2,7\},\ \{3,7\},\ \{3,6\},\ \{3,5\},\ \{0,3\},\text{ and }\{0,2\}.$$

Hence, any minimal path between W_8^- and W_8^+ visits at most 7 triangulations. It follows that the flip distance of A_8 is precisely equal to 7. ☐

When $9 \leq n \leq 12$, the flip distance of A_n can be found using proposition 7. Observe that the lower bound on $\delta(A_n)$ stated by this proposition depends on $\delta(B_{n-1})$ and $\delta(C_{n-1})$. Recall that B_n is defined for every n greater than 7 as the unordered pair that contains triangulations Y_n^- and Y_n^+, and that C_n is obtained for the same values of n replacing Y_n^- by X_n^- in B_n. Triangulations Y_n^-, X_n^-, and Y_n^+ are shown in figure 10 when $8 \leq n \leq 11$. The edges of triangulation Y_n^- are drawn as solid lines in the top of this figure. Triangulation X_n^- is obtained from Y_n^- replacing edge $\{3,6\}$ by edge $\{4,5\}$, depicted as a dotted line.

Now observe that, for any $T \in \{Y_8^-, X_8^-\}$, triangulation Y_8^+ can be flipped to T using a sequence of flips so that each flip carried out in this sequence introduces an edge of T. Hence, it follows from lemma 2 that $\delta(\{Y_8^+,T\})$ is equal to the number of interior edges in T, which proves that $\delta(B_8)$ and $\delta(C_8)$ are both equal to 5. The flip distances of B_n and C_n can further be obtained easily when $9 \leq n \leq 11$ using the results proven in the previous sections:

Proposition 9 *Let n be an integer. If $8 \leq n \leq 11$, then $\delta(B_n)$ and $\delta(C_n)$ are both equal to $2n-11$.*

Proof First, if n is equal to 8, then $2n-11$ is equal to 5. Hence, in this case, the desired equalities immediately follow from the discussion that precedes the proposition. Now assume that $9 \leq n \leq 11$ and consider a pair $P \in \{B_n, C_n\}$.



Fig. 10 Triangulations Y_n^- and Y_n^+, depicted in solid lines when n ranges from 8 to 11. Triangulations X_n^-, obtained by flipping edge $\{4, 6\}$ in Y_n^- are sketched using dotted lines for the same values of n.

The triangulation in P that is distinct from Y_n^+ will be called T. Hence, T is either equal to Y_n^- or to X_n^-. Denote:

$$\upsilon = \begin{cases} \{1, 3\} \text{ if } 9 \leq n \leq 10, \\ \{0, 3\} \text{ if } n = 11. \end{cases}$$

Further call $Q = \Lambda_\upsilon^\star(P)$. It can be seen in figure 10 that the links of edges $\{1, 2\}$ and $\{2, 3\}$ in T are always respectively $\{7\}$ and $\{6\}$ and that the link of $\{1, 3\}$ in Y_n^+ is always $\{0, 2\}$. As these three links are disjoint, and as $\{1, 2, 3\}$ is disjoint from $\{6, 7\}$, then according to theorem 6, $\delta(P) \geq \delta(Q) + 3$ when $9 \leq n \leq 10$. Now observe that, if $n = 11$, then the link of $\{0, 1\}$ in T is $\{8\}$ and the link of $\{0, 3\}$ in Y_n^+ is $\{1, 10\}$. As a consequence, for this value of n, the links of $\{0, 1\}$, $\{1, 2\}$, and $\{2, 3\}$ in T and that of $\{0, 3\}$ in Y_n^+ are pairwise disjoint. In addition, $\{6, 7, 8\}$ and $\{0, 1, 2, 3\}$ are disjoint. Hence, theorem 6 yields $\delta(P) \geq \delta(Q) + 5$ when $n = 11$.

Denote $k = 3$ if $9 \leq n \leq 10$ and $k = 5$ if $n = 11$. It has been shown is the last paragraph that the following inequality holds:

$$\delta(P) \geq \delta(Q) + k. \tag{20}$$

Now consider pair Q. One can see that the two triangulations in this pair do not have any interior edge in common. As a consequence, the flip distance of Q is not less than the number of interior edge in the triangulations it contains. In particular, when $n = 9$, the flip distance of Q is not less than 4. Further observe that, when $10 \leq n \leq 11$, the edge introduced in $\Lambda_I^\star(Y_n^+)$ by any possible flip never belongs to $\Lambda_I^\star(T)$. Hence, for these values of n, the flip distance of Q is greater than the number of interior edges in the triangulations it contains. As these triangulations have 5 interior edges when $10 \leq n \leq 11$, then the flip distance of Q is not less than 6 for these values of n.

Denote $l = 4$ if $n = 9$ and $l = 6$ if $10 \leq n \leq 11$. It has been shown is the last paragraph that the following inequality holds:

$$\delta(Q) \geq l. \tag{21}$$

Combining inequalities (20) and (21), one obtains that $\delta(P) \geq 2n-11$ when $9 \leq n \leq 11$. The opposite inequality is found by exhibiting an adequate path between the two triangulations in pair P. □

Invoking propositions 7, 8, and 9, one finds that the flip distance of A_n is never less than $2n-10$ when $3 \leq n \leq 12$:

Lemma 11 *Let n be an integer. If $3 \leq n \leq 12$, then $\delta(A_n) \geq 2n-10$.*

Proof The lemma is proven by induction on n. If $3 \leq n \leq 8$, then the result immediately follows from proposition 8. Now assume that for some integer n so that $8 < n \leq 12$, and for any $i \in \iota(n)$, the flip distance of pair A_i is not less than $2i-10$. As n is greater than 8, proposition 7 yields:

$$\delta(A_n) \geq \min(\delta(A_{n-1})+2, \delta(A_{n-2})+4, \delta(B_{n-1})+3, \delta(C_{n-1})+3). \tag{22}$$

By induction, the flip distances of A_{n-1} and A_{n-2} are not less than respectively $2n-12$, $2n-14$. In addition, since $8 \leq n-1 \leq 11$, proposition 9 states that $\delta(B_{n-1})$ and $\delta(C_{n-1})$ are both equal to $2n-13$. Hence, the result follows from inequality (22). □

Lemma 11 gives a lower bound on the flip distance of A_n when $3 \leq n \leq 12$. Using the recursive expression stated by theorem 12, one can generalize this lower bound to arbitrarily large integers n:

Theorem 13 *For any integer n greater than 2, $\delta(A_n) \geq 2n-10$.*

Proof The theorem is proven by induction on n. If $n \leq 12$, then the result immediately follows from lemma 11. Now assume that for some integer $n > 12$, and for any $i \in \iota(n)$, the flip distance of pair A_i is not less than $2i-10$. As n is greater than 12, theorem 12 yields:

$$\delta(A_n) \geq \min(\delta(A_{n-1})+2, \delta(A_{n-2})+4, \delta(A_{n-5})+10, \delta(A_{n-6})+12). \tag{23}$$

By induction, the flip distances of A_{n-1}, A_{n-2}, A_{n-5}, and A_{n-6} are not less than respectively $2n-12$, $2n-14$, $2n-20$, and $2n-22$. Combining these four inequalities with (23) completes the proof. □

It follows from lemma 1 and theorem 13 that the flip distance of A_n is precisely $2n-10$ when n is greater than 12. According to theorem 2, the diameter of associahedra with dimension above 9 is found as a consequence:

Corollary 4 *For any integer d greater than 9, the d-dimensional associahedron has diameter $2d-4$.*

9 Discussion

It has been shown using combinatorial arguments that triangulations W_n^- and W_n^+ have flip distance $2n-10$ when n is greater than 12, thus settling the two open problems on the diameters of associahedra [17,4,6,5]. It is first shown in this concluding section that, in fact, these triangulations correspond to maximally distant vertices of the d-dimensional associahedron whenever d is distinct from 6, 7, and 9. A third related open problem, originally posed in [4], is solved next. A discussion on the possible extension of the techniques developed in sections 3 and 4 to a class of polytopes that generalize associahedra completes the section.

9.1 The diameters of d-dimensional associahedra when $d \leq 9$

Combining lemma 1 with theorems 2 and 13, one obtains that triangulations W_{d+3}^- and W_{d+3}^+ correspond to maximally distant vertices of the d-dimensional associahedron when d is greater than 9. When d is not greater than 9, the diameters of associahedra can be obtained using a computer program. These diameters are respectively 0, 1, 2, 4, 5, 7, 9, 11, 12, and 15 when d ranges from 0 to 9 (see [17]). Hence, according to proposition 8 and to lemma 11, W_{d+3}^- and W_{d+3}^+ further correspond to maximally distant vertices of the d-dimensional associahedron when d is either less than 6 or equal to 8. It is worth noting that this property fails in the case of the three remaining dimensions, i.e. when d is equal to 6, 7, or 9. Indeed, recall that the flip distance of any two triangulations of $\iota(n)$ is not greater than $2n-10$ when n is greater than 12. In the case of triangulations W_n^- and W_n^+, this observation remains true down to $n=9$, which can be shown using a proof similar to that of lemma 2. in [17]:

Proposition 10 *For any integer n greater than 8, $\delta(A_n) \leq 2n-10$.*

Proof Let n be an integer greater than 8. Observe that for these values of n, vertex 0 is always contained in exactly four interior edges of W_n^+. Further note that it is always possible to find a flip that increases the number of interior edges incident to some vertex in a given triangulation. As W_n^+ has $n-3$ interior edges, then one can transform W_n^+ into the unique triangulation T of $\iota(n)$ all of whose interior edges admit 0 as a vertex by carrying out $n-7$ flips. For the same reason, one can transform W_n^- into T with a sequence of $n-3$ flips. This shows that the flip distance of W_n^- and W_n^+ is at most $2n-10$. □

It follows from this proposition that $\delta(A_{d+3})$ is less by one than the diameter of the d-dimensional associahedron when $d \in \{6,7,9\}$.

9.2 A third open problem solved

A conjecture from [4] is now disproved. Consider an integer n greater than 3, and observe that $\{n-1,n+1\}$ and $\{0,2\}$ are two edges on $\iota(n+2)$. One



Fig. 11 Triangulations M_n^- (left) and M_n^+ (right), depicted when n is greater than 5. The interrupted edges show that the length of each zigzag is arbitrary.

can define two triangulations of $\iota(n)$ by respectively contracting these edges in $Z_{n+2}^-(0)$ and in $Z_{n+2}^+(-1)$:

$$M_n^- = \Lambda^{\star}_{\{n-1,n+1\}}(Z_{n+2}^-(0)) \text{ and } M_n^+ = \Lambda^{\star}_{\{0,2\}}(Z_{n+2}^+(-1)).$$

These two triangulation are shown in figure 11, depending on the parity of n. Each of them admits a unique comb with three teeth. Observe that M_n^- and M_n^+ could have been alternatively defined by carrying out a flip in respectively $Z_n^-(1)$ and $Z_n^+(-2)$. Consider the following pair of triangulations:

$$D_n = \{M_n^-, M_n^+\}.$$

It is conjectured in [4] that D_n has flip distance $2n-10$ when n is greater than 9. Observe that this conjecture was stated in the formalism of binary trees (see conjecture 5.1. in [4]). While this conjecture turns out to be false, the insight provided in [4] on the form of two possibly maximally-distant vertices of the associahedra was partly correct. Indeed, pairs A_n and D_n are similar, except for the facts that the two triangulations in A_n are not symmetric when n is even and have a comb at each end of the zigzag. The two triangulations in pair D_n are not always at maximal distance, though:

Proposition 11 *For any integer n greater than* 19, $\delta(D_n) \leq \delta(D_{n-16}) + 31$.

Proof Let n be an integer greater than 19. Because of this condition on n, the following fourteen edges are interior edges of triangulation M_n^-:

$$\begin{array}{l}\{2,n-1\},\ \{1,n-1\},\ \ \{4,n-2\},\ \{4,n-3\},\ \{5,n-3\},\\ \{5,n-4\},\ \{6,n-4\},\ \ \{7,n-5\},\ \{6,n-5\},\ \{7,n-6\},\\ \{8,n-6\},\ \{10,n-8\},\ \{9,n-8\},\ \{9,n-7\}.\end{array}$$

Let ψ be the path of length 14 that starts at triangulation M_n^- and that successively flips these edges in the order in which they are listed above, from left to right and then from top to bottom. Similarly, since n is greater than 19, the following seventeen edges are interior edges of M_n^+:

$$\begin{array}{l}\{2,n-6\},\ \{2,n-5\},\ \ \{1,n-5\},\ \ \{1,n-4\},\ \ \{0,n-4\},\ \{0,n-3\},\\ \{0,n-2\},\ \{4,n-8\},\ \ \{4,n-7\},\ \ \{5,n-8\},\ \ \{5,n-9\},\ \{5,n-7\},\\ \{3,n-7\},\ \{7,n-11\},\ \{7,n-10\},\ \{6,n-10\},\ \{6,n-9\}.\end{array}$$



Fig. 12 Pairs $\{\psi(7), \phi(7)\}$ (left), $\{\psi(11), \phi(13)\}$ (center), and $\{\psi(14), \psi(17)\}$ (right) defined in the proof of proposition 11. The edges that belong to the two triangulations of a given pair, and whose links in these two triangulations are equal, are drawn using dotted lines.

Call ϕ the path of length 17 that starts at triangulation M_n^+ and that successively flips these edges in the order in which they are listed above, from left to right and then from top to bottom.

In order to visualize paths ψ and ϕ, pairs $\{\psi(7), \phi(7)\}$, $\{\psi(11), \phi(13)\}$, and $\{\psi(14), \psi(17)\}$ are respectively sketched in the left, in the center and in the right of figure 12. It can be seen that each of the first seven flips along these two paths introduces an edge that admits 3 as a vertex. In addition, $\psi(7)$ and $\phi(7)$ both admit a triangulation of $\iota(n-4) \setminus \iota(3)$ as a subset, drawn using solid lines. The edges of $\psi(7)$ and $\phi(7)$ that are not edges on $\iota(n-4) \setminus \iota(3)$ are further shown as dotted lines, and it can be seen that they all belong to both $\psi(7)$ and $\phi(7)$. By construction, none of these edges are removed along paths ψ and ϕ. Hence, they are contained in all the subsequent triangulations visited by paths ψ and ϕ, and for this reason, they are omitted from the sketches in the central and rightmost columns in figure 12.

Similarly, $\psi(11)$ and $\phi(13)$ each admit a triangulation of $\iota(n-6) \setminus \iota(6)$ as a subset, depicted using solid lines in the center of figure 12. Moreover, the edges of $\psi(11)$ and $\phi(13)$ that are not edges on $\iota(n-6) \setminus \iota(6)$ are common to $\psi(11)$ and $\phi(13)$ and are not removed along paths ψ and ϕ. These edges are omitted from the sketches of $\psi(14)$ and $\phi(17)$.

Finally, it can be seen that $\psi(14)$ and $\phi(17)$ each contain a triangulation of $\iota(n-8) \setminus \iota(8)$ as a subset, depicted using solid lines in the right of figure 12. In fact, these triangulations are the respective images of M_{n-16}^- and M_{n-16}^+ by the following boundary-preserving bijection:

$$\alpha : \begin{array}{l} \pi \to \iota(n-8) \setminus \iota(8), \\ i \mapsto i + 8, \end{array}$$

where $\pi = \iota(n-16)$. Moreover, all the edges of $\psi(7)$ and $\phi(7)$ that are not edges on $\iota(n-8) \setminus \iota(8)$ belong to both $\psi(14)$ and $\phi(17)$. Hence, it follows from lemma 6 that:

$$\delta(\{\psi(14), \phi(17)\}) = \delta(D_{n-16}).$$

As ψ is a path of length 14 from M_n^- to $\psi(14)$ and as ϕ is a path of length 17 from M_n^+ to $\psi(17)$, then one obtains $\delta(D_n) \leq \delta(D_{n-16}) + 14 + 17$ which completes the proof. $\square$

According to this proposition, there exists an integer k so that:

$$\delta(D_n) \leq \frac{31}{16}n + k.$$

Hence, $\delta(D_n)$ is not only less than $\delta(A_n)$ when n is large enough, but the difference of these two quantities increases at least linearly with n.

9.3 On the diameters of more general polytopes

Along with the many constructions [1,2,3,6,9,14,15] that followed the ones by Mark Haiman and Carl Lee, several generalizations of associahedra have been discovered [1,2,8,9] in the last twenty years. Among them, one finds the secondary polytopes [9,1] and the fiber polytopes [2]. Many properties of associahedra carry over to these polytopes. In particular, their vertices are in one-to-one correspondence with certain triangulations of given finite sets of points. It is then natural to ask whether the techniques developed in this article could also be generalized to such triangulations, in the hope for results on the diameters of these more general polytopes.

Recall that contracting an edge in some triangulation of a polygon always results in a triangulation. This is not true any more in the case of more general triangulations, even already in dimension 2 for triangulations with interior vertices. In fact, the notion of contraction itself is not so easy to define for more general triangulations because their vertices cannot, in general, be ordered. A possible starting point would be to exclusively consider the contractions of single boundary faces. Even in this case, though, the resulting objects are not necessarily triangulations. In other words, not every boundary face of a triangulation could be contracted. Another possible starting point would be to consider triangulations of sets of points with good properties, such as the vertex sets of cyclic polytopes. These polytopes beautifully generalize polygons in higher dimensions. In particular, their vertices can be identified with points on the moment curve [10], and can therefore be ordered as these points. It has been shown that contracting successive vertices with respect to this ordering in a triangulation of a cyclic polytope always results in a triangulation [16]. Hence, the techniques developed in this article may be applied to such point configurations in the search for results on the diameters of their secondary or fiber polytopes. Note however that, unlike associahedra, the number of vertices of these polytopes is unknown in general, and that no reasonable upper bound is available on their diameters.

References

1. L. J. Billera, P. Filliman, B. Sturmfels, Construction and complexity of secondary polytopes, Adv. Math. **83**, 155–179 (1990)
2. L. J. Billera, B. Sturmfels, Fiber polytopes, Ann. Math. **135**, 527-549 (1992)
3. F. Chapoton, S. Fomin, A. Zelevinsky, Polytopal realizations of generalized associahedra, Canad. Math. Bull. **45**, 537–566 (2002)
4. P. Dehornoy, On the rotation distance between binary trees, Adv. Math. **223**, 1, 1316–1355 (2010)
5. J. A. de Loera, J. Rambau, F. Santos, Triangulations: structures for algorithms and applications, Algorithms and Computation in Mathematics 25, Springer (2010)
6. S. L. Devadoss, A realization of graph associahedra, Discrete Math. **309**, 271–276 (2009)
7. S. L. Devadoss, J. O'Rourke, Discrete and Computational Geometry, Princeton University Press, 2011.
8. S. Fomin, A. Zelevinsky, Y-systems and generalized associahedra, Ann. Math. **158**, 977–1018 (2003)
9. I. M. Gel'fand, M. M. Kapranov, A. V. Zelevinsky, Discriminants, resultants, and multidimensional determinants, in: Mathematics: Theory & Applications, Birkhäuser, Boston, 1994
10. B. Grünbaum, Convex Polytopes, Graduate texts in mathematics **221**, Springer (2003)
11. M. Haiman, Constructing the associahedron, unpublished manuscript, MIT (1984)
12. D. Huguet, D. Tamari, La structure polyédrale des complexes de parenthésages, J. Comb. Inf. Sys. Sci. **3**, 69–81 (1978)
13. C. W. Lee, The associahedron and triangulations of the n-gon, Eur. J. Comb. **10**, 551–560 (1989)
14. J.-L. Loday, Realization of the Stasheff polytope, Arch. Math. **83**, 267–278 (2004)
15. A. Postnikov, Permutohedra, associahedra, and beyond, Int. Math. Res. Notices **2009**, 6, 1026–1106 (2009)
16. J. Rambau, F. Santos, The Generalized Baues Problem for Cyclic Polytopes I, Eur. J. Comb. **21**, 65–83 (2000)
17. D. Sleator, R. Tarjan, W. Thurston, Rotation distance, triangulations, and hyperbolic geometry, J. Amer. Math. Soc. **1**, 647–681 (1988)
18. J. Stasheff, Homotopy associativity of H-spaces I, Trans. Amer. Math. Soc. **108**, 275–292 (1963)